

BEHAVIORAL RISK FACTOR SURVEY TABLES, AGES 18 AND OLDER

	2009	2008	2007	2006	2005	2004	2003	2002	2001
Estimated Alcohol Use in Past 30 Days (%)									
Missouri	50.1	51.9	51.9	50.7	51.8	51.8	53.5	53.6	49.6
U.S. (Median of the States)	54.3	54.4	54.8	55.4	56.2	57.1	59.4	58.1	55.8
Estimated Binge Drinking * in Past 30 Days (%)									
Missouri	17.2	15.3	16.2	16.5					
U.S. (Median of the States)	15.7	15.6	15.8	15.4					
* 5 or more drinks on one occasion for males									
* 4 or more drinks on one occasion for females									
Heavy Drinking ** in Past 30 Days (%)									
Missouri	4.1	5.3	4.6	6.2	5.0	6.0	5.5	6.1	4.8
U.S. (Median of the States)	5.1	5.1	5.2	4.9	4.9	4.9	5.8	5.9	5.1
** More than 2 drinks per day for males									
** More than 1 drink per day for females									
Estimated Tobacco Smoking in Past 30 Days (%)									
Missouri	23.1	24.9	24.5	23.2	23.4	24.1	27.2	26.5	25.9
U.S. (Median of the States)	17.9	18.3	19.8	20.1	20.6	20.9	22.0	23.2	23.2
Estimated Former Smokers (%)									
Missouri	24.7	25.0	23.4	26.8	24.2	24.2	24.3	25.5	24.5
U.S. (Median of the States)	25.5	25.2	24.6	24.7	24.8	24.1	25.2	24.7	24.7

Values shown are estimates based on survey samples. Apparent differences in values between years, and between U.S. and Missouri, may not be statistically significant.

Data Source:

U.S. Department of Health and Human Services, Centers for Disease Control and Prevention: Behavioral Risk Factor Surveillance System