



# Behavioral Health Profile for Nodaway County



July 1, 2011

## About Nodaway County:

Located in Northwest Missouri, Nodaway County has a population of 23,370. Among Missouri's 114 counties plus the city of St. Louis, Nodaway County ranks 49th in terms of population. Nodaway County's unemployment rate in 2010 was 6.64%. This was lower than the statewide unemployment rate of 9.60%. In 2009, the poverty rate for Nodaway County was 21%. This was higher than the statewide poverty rate of 14.6%. Nodaway County is not considered a primary healthcare shortage area and is not considered a mental health professional shortage area by the Health Resources and Services Administration. Other shortages may exist for low income populations.

## Substance Use and Abuse in Nodaway County:

The availability of county-level data on substance use and abuse is limited. The Missouri Student Survey can provide estimates for youth in most Missouri counties. This survey is administered in even-numbered years to 6th through 12th grade students in participating school districts.

In Nodaway County, 52.54% of youth believe that it would be easy to get cigarettes and 39.63% have friends who smoke. The Healthy People 2020 has established objectives to reduce tobacco initiation and use by adolescents and adults. An estimated 62.26% youth in Nodaway County believe that it would be easy to get alcohol and 58.57% have friends who drink alcohol. Youth who use alcohol are at greater risk of developing abuse problems later in life. In Nodaway County, 22.94% of youth believe that it would be easy to get marijuana and 9.94% of youth believe that it would be easy to get other drugs such as cocaine, methamphetamine, and ecstasy. Missouri youth tend to perceive marijuana as being safer than other illegal drugs and safer than tobacco.

In Northwest Missouri, an estimated 7.74% of residents age 12 and over use illegal drugs and 7.98% are abusing alcohol. Public safety, health care, crime, and public assistance are among the areas impacted by substance abuse. In 2009, Nodaway County had 27 alcohol-related traffic crashes. This is higher than that for 2008 (23). Of the 2009 alcohol-related crashes, 4 crashes resulted in the loss of life. Another 12 resulted in injuries. Alcohol-related crashes are more likely to produce fatalities and injuries compared to non-alcohol-related crashes.

In 2009, Nodaway County had 166 DUI arrests, 231 liquor law violations, and 80 drug arrests. Nodaway County had 4 methamphetamine laboratory seizures in 2009. The manufacture of methamphetamine has been particularly problematic for rural Missouri.

## Current Substance Use for Grades 6 through 12\*


	Nodaway County	Missouri
Cigarettes	8.32 %	13.70 %
Chewing tobacco	<b>7.58 %</b>	6.10 %
Alcohol	14.20 %	19.80 %
Binge drinking**	10.26 %	11.40 %
Marijuana	1.72 %	9.40 %
Inhalants	1.14 %	3.00 %
Prescription drugs not prescribed for student	3.61 %	6.70 %
Over-the-counter medications for non-medical use	2.46 %	4.80 %

\*Used in the past 30 days

**Red:** Higher than Missouri rate

\*\*Five or more drinks on one occasion

## 2009 Alcohol-related Crashes




# Behavioral Health Profile for Nodaway County

## Substance Use and Abuse in Nodaway County: (cont.)

In 2008, Nodaway County residents had a total of 112 alcohol-related and 95 drug-related hospitalizations and emergency room visits. Statewide, average expenses for an alcohol or drug-related visit to the emergency room was \$4,180.

In state fiscal year 2010, 73 Nodaway County residents were admitted to substance abuse treatment at publicly-funded facilities. In Northwest Missouri and statewide, alcohol and marijuana were the most common substance abuse problems presenting for treatment. In Northwest Missouri, the next most common drug problem was methamphetamine. Statewide, it was methamphetamine.



## Mental Health in Nodaway County:

In Northwest Missouri, an estimated 13.97% of residents age 12 and over have had serious psychological distress in the past year. Young adults and females tend to have higher rates of serious psychological distress. Nationwide, about 45 percent of those with serious psychological distress receive mental health services. Approximately 8.66% of Northwest Missouri residents have had at least one major depressive episode in the past year. A major depressive episode is characterized by an extended period of depressed mood, loss of interest or pleasure, and impaired functioning. Females are more likely to report having had a major depressive episode. Mental illness can result from a combination of genetic vulnerabilities and life stressors such as loss of employment or relationships and financial difficulties. Symptoms can include feeling sad, confused thinking, excessive fears, withdrawal from family and friends, problems sleeping, detachment from reality, substance abuse, significant changes in eating habits, excessive anger or violence, and suicidal thinking. Treatment for mental illness can include a combination of counseling and medication. In 2009, 5 Nodaway County residents committed suicide. The suicide rate for Northwest Missouri was 14.1 per 100,000 persons, which is lower than the state rate (14.3 per 100,000 persons.) Males are about four times more likely to commit suicide than females. Older males have higher rates of suicide than younger males.

In state fiscal year 2010, 132 Nodaway County residents received treatment for serious mental illness at publicly-funded facilities. In Northwest Missouri and statewide, mood disorders were the most common diagnosis category. Mood disorders include mania, major depression, and bipolar disorder. Anxiety and psychotic disorders were the next most common diagnosis categories. Anxiety disorders include panic, obsessive-compulsive, and post-traumatic stress disorders as well as phobias. Psychotic disorders include schizophrenia and delusional disorders. Individuals struggling with serious mental illness are at higher risk for homicide, suicide, and accidents as well as chronic conditions including cardiovascular and respiratory diseases and substance abuse disorders.

	Nodaway County	Missouri
Were very sad	16.64 %	18.90 %
Were grouchy, irritable, or in a bad mood	<b>25.33 %</b>	24.80 %
Felt hopeless about the future	11.21 %	11.80 %
Felt like not eating or eating more than usual	15.43 %	18.10 %
Slept a lot more or a lot less than usual	20.11 %	25.60 %
Had difficulty concentrating on school work	22.24 %	24.60 %
Seriously considered attempting suicide	11.01 %	11.60 %
Have been bullied on school property	<b>25.61 %</b>	25.00 %

\*In the past 30 days

**Red:** Higher than Missouri rate