



# Behavioral Health Profile

## Wayne County


2018



Wayne County is located in Southeast Missouri and had a population of 13,207 in 2016. Wayne County ranks 77 in population size among Missouri's 114 counties plus the city of St. Louis. The 2016 unemployment rate in the county was 9.1%. This was higher than the statewide unemployment rate of 6.6%. In 2016, the poverty rate was 26.0% which was higher than the statewide poverty rate of 14.0%. The median income of the county was \$30,793.

### Substance Use in Wayne County

The availability of county-level data on substance use is limited. The Missouri Student Survey convenience sample can provide estimates for youth in most Missouri counties. This survey is administered in even-numbered years to 6th through 12th grade students in participating school districts. Adult data are available at the regional level from the NSDUH. See page 2 for graph.




**Alcohol** is one of the most common substances used by Missouri youth. In Wayne County, 16.0% youth have used alcohol in the past 30 days. An estimated 35.1% believe that using alcohol presents only 'slight' or 'no risk' of harm. Similarly, approximately 32.2% of youth believe that there is 'slight' or 'no risk' of harm in binge drinking once or twice a week. About 52.0% of youth believe that it would be easy to get alcohol. 40.9% of youth have at least one friend that uses alcohol.

The average age of first alcohol use is 12.3. Among Missouri adults, alcohol is the most commonly used substance. In the Southeast region, approximately 47.2% of adults currently drink alcohol, and 25.8% have had 5 or more drinks of alcohol on a single occasion in the past 30 days.

**Cigarette** use, in both standard and electronic forms, is of concern across the state. In Wayne County youth, the rate of e-cigarette use is higher than the rate of standard cigarette use. The average age that county youth first use standard cigarettes is 11.8. An estimated 56.4% of youth believe that it would be easy to get standard cigarettes and 50.2% to get e-cigarettes. 36.4% of youth have at least one friend that uses standard cigarettes. An estimated 15.1% of youth believe there is 'slight' or 'no risk' to smoking standard cigarettes, while 40.7% believe there is 'slight' or 'no risk' to smoking e-cigarettes. Most youth obtain cigarettes and e-cigarettes from their friends. In the Southeast region, 34.3% of adults currently use cigarettes compared to 29% statewide.

**Marijuana** use in Missouri youth continues to be of interest, particularly with recent shifts in legality of adult use across the nation. Around 7.5% of youth and 7.3% of adults report current marijuana use. In Wayne County, 5.1% of youth use marijuana. In 2018, 40.6% of youth said that marijuana is easy to get. Around 80.2% believe that smoking marijuana is wrong, and 33.7% report having at least one friend that uses marijuana. The average age of first use is 13.5. The majority of youth obtain marijuana from their friends, a dealer, or a family member. Across the state, around 46.6% of adults believe there is 'slight' or 'no risk' of harm from smoking marijuana once or twice a week. In Wayne County youth, approximately 35.4% believe that there is 'slight' or 'no risk' of harm from smoking marijuana once or twice a week.

**Prescription drug** misuse is of growing concern both across the nation and in Missouri. In Wayne County, 31.3% of youth believe that it would be easy to get prescription drugs that were not prescribed to them. Most youth report that there is a risk of harm when misusing prescription drugs, but 11.5% of youth believe that there is 'slight' or 'no risk at all'. About 95.7% of youth believe it would be wrong to use prescription drugs not prescribed to them. Most youth get prescription drugs from a friend or family member. Pain medication is the most commonly misused prescription medication, with 44.4% of those misusing prescription drugs in Wayne County county saying they misuse pain medication. 4.0% of adults in the Southeast region reported misuse of prescription pain medication over the past year.



### Substance Use Consequences in the County

**Health:** In 2015, Wayne County residents had a total of 6 alcohol-related and 11 drug-related hospitalizations. In addition there were 11 alcohol-related and 21 drug-related ER visits that did not include a hospital stay.

**Treatment:** In 2017, 104 individuals were admitted into Substance Abuse Treatment Programs. A total of 23 were primarily due to alcohol, 21 were primarily due to marijuana, and 8 were primarily due to prescription drugs.

**Law Enforcement:** In 2017, Wayne County had 37 DWI arrests, 5 liquor law violations and 146 drug-related arrests. There were 1 methamphetamine laboratory seizures in Wayne County in 2017.

**Traffic Crashes:** Alcohol-related traffic crashes decreased in the last year (from 21 in 2015 to 20 in 2016). Alcohol-related crashes are more likely to produce fatalities and injuries compared to non-alcohol-related crashes.

2016 Number of Alcohol-Related Crashes by Severity				2016 Number of People Injured / Killed in Alcohol and Drug-Related Crashes			
Total Crashes	Fatal Crash	Crash w/ Injury	Crash w/ Property Damage	Alcohol Fatalities	Alcohol Injuries	Drug Fatalities	Drug Injuries
20	0	12	8	0	18	1	9



### Mental Health Data for Wayne County

Individuals struggling with serious mental illness are at higher risk for homicide, suicide, and accidents, as well as chronic conditions including cardiovascular disease, respiratory disease, and substance use disorders. In state fiscal year 2017, 257 Wayne County residents received treatment for serious mental illness at publicly-funded facilities. See table for details. While there are data on those who receive treatment, data on mental health in the general population is very limited. This is especially true at the local level.

Numbers Served in Wayne County		
	FY2016	FY2017
Total	225	257
Adjustment Disorder	*	*
Anxiety Disorder	39	37
Developmental Disorder	0	0
Impulse Control Disorder	9	12
Mood Disorder	140	125
Psychotic Disorder	32	26

These numbers indicate the number of clients seen with each diagnosis per year. An individual client may have more than one admission within a year.

\*Counts under 5 suppressed to protect identities

In Southeast Missouri, 20.7% of those 18 and older had a mental illness in the past year, and 6.1% had a serious mental illness. Serious mental illness is defined as any of the mental disorders asked about and 'these disorders resulted in substantial impairment in carrying out major life activities'.

Approximately 7.5% of Southeast Missouri residents ages 18+ had at least one major depressive episode in the past year. A major depressive episode is characterized by an extended period of depressed mood, loss of interest or pleasure, and impaired functioning. Typically, females are more likely to report having had a major depressive episode.

Students (6th-12th grade) in the county were asked about their mental health. 74.9% of students in the county reported feeling grouchy or in a bad mood, 59.5% reported feeling sad, 56.7% had changes in sleep, and 62.3% had difficulty concentrating in school at least sometimes. About 19.5% of youth reported self-harm. The most common method of self-harm was cutting, hitting, or scratching oneself. Females were more likely to report self-harm than males.

Suicide is the 2nd leading cause of death for ages 10-34 in Missouri. In 2016, 5 Wayne County residents died by suicide. Typically, white males are most at risk of suicide. Approximately 10.7% of youth had considered suicide in the last year, 9.6% made a plan, and 1.1% actually attempted, resulting in an injury.

Developed with support from the Missouri Division of Behavioral Health and the Substance Abuse and Mental Health Services Administration.

For more information: <http://dmh.mo.gov/ada/mobhew/>