

# UNITED STATES DRUG USE: AGE 12 AND OLDER

## National Survey on Drug Use and Health

	2004	2003	2002	2001	2000	1999	1998	1997	1996	1995	1994	1993	1992
<b>PAST 30-DAY USE (%)</b>													
Alcohol	50.3	50.1	51.0	48.3	46.6	46.4	51.7	51.4	51.0	52.2	53.9	50.8	49.0
Binge Alcohol Use *	22.8	22.6	22.9	20.5	20.6	20.2	15.6	15.3	15.5	15.8	16.5	14.6	14.5
Heavy Alcohol Use **	6.9	6.8	6.7	5.7	5.6	5.7	5.9	5.4	5.4				
Any Tobacco	29.2	29.8	30.4	29.5	29.3	30.2							
Cigarettes	24.9	25.4	26.0	24.9	24.9	25.8	27.7	29.6	28.9	28.8	28.6	29.6	31.9
Smokeless Tobacco	3.0	3.3	3.3	3.2	3.4	3.4	3.1	3.2	3.2	3.3	3.3	3.2	4.0
Any Illicit Drug	7.9	8.2	8.3	7.1	6.3	6.3	6.2	6.4	6.1	6.1	6.0	5.9	5.8
Marijuana/Hashish	6.1	6.2	6.2	5.4	4.8	4.7	5.0	5.1	4.7	4.7	4.8	4.6	4.7
Inhalant	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.4	0.3	0.3
Hallucinogen	0.4	0.4	0.5	0.6	0.4	0.4	0.7	0.8	0.6	0.7	0.5	0.4	0.4
LSD	0.1	0.1	0.0	0.1	0.2	0.2	0.3	0.2	0.2	0.3	0.2		
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0
Ecstasy	0.2	0.2	0.3	0.3									
Cocaine (any form)	0.8	1.0	0.9	0.7	0.5	0.7	0.8	0.7	0.8	0.7	0.7	0.7	0.7
Crack Cocaine	0.2	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2
Heroin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.0
Pain Reliever (non-medical use)	1.8	2.0	1.9	1.6	1.2	1.2	0.8	0.7	0.9	0.6	0.7	0.8	0.9
Stimulant (non-medical use)	0.5	0.5	0.5	0.5	0.4	0.4	0.3	0.3	0.4	0.4	0.3	0.5	0.3
Methamphetamine	0.2	0.3	0.3	0.3	0.2	0.2							
Sedative (non-medical use)	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.2
Tranquilizer (non-medical use)	0.7	0.8	0.8	0.6	0.4	0.5	0.3	0.4	0.4	0.4	0.5	0.6	0.8
<b>LIFETIME USE (%)</b>													
Alcohol	82.4	83.1	83.1	81.7	81.0	81.3	81.3	81.9	82.6	82.3	84.2	82.6	81.9
Any Tobacco	71.4	72.7	73.1	71.4	70.5	72.0							
Cigarettes	67.3	68.7	69.1	67.2	66.5	68.2	69.7	70.5	71.6	71.8	73.3	73.3	73.1
Smokeless Tobacco	18.6	19.4	19.9	19.1	18.5	19.1	17.2	17.3	17.0	17.0	17.2	15.9	18.2
Any Illicit Drug	45.8	46.4	46.0	41.7	38.9	39.7	35.8	35.6	34.8	34.2	34.4	34.2	33.3
Marijuana/Hashish	40.2	40.6	40.4	36.9	34.2	34.6	33.0	32.9	32.0	31.0	31.1	31.0	30.2
Inhalant	9.5	9.7	9.7	8.1	7.5	7.8	5.8	5.7	5.6	5.7	5.8	5.9	5.3
Hallucinogen	14.3	14.5	14.6	12.5	11.7	11.3	9.9	9.6	9.7	9.5	8.7	9.0	8.3
LSD	9.7	10.3	10.4	9.0	8.8	8.7	7.9	7.8	7.7	7.5	7.0	7.0	6.7
PCP	2.8	3.0	3.2	2.7	2.6	2.6	3.5	3.0	3.2	3.2	2.8	2.7	2.7
Ecstasy	4.6	4.6	4.3	3.6	2.9								
Cocaine (any form)	14.2	14.7	14.4	12.3	11.2	11.5	10.6	10.5	10.3	10.3	10.4	11.3	10.9
Crack Cocaine	3.3	3.3	3.6	2.8	2.4	2.7	2.0	1.9	2.2	1.8	1.9	1.9	1.5
Heroin	1.3	1.6	1.6	1.4	1.2	1.4	1.1	0.9	1.1	1.2	1.0	1.0	0.8
Pain Reliever (non-medical use)	13.2	13.1	12.6	9.8	8.6	9	5.3	4.9	5.5	6.1	6.0	6.4	6.1
Stimulant (non-medical use)	8.3	8.8	9.0	7.1	6.6	7.2	4.4	4.5	4.7	4.9	4.6	4.8	5.0
Methamphetamine	4.9	5.2	5.3	4.3	4.0	4.3	2.1	2.5	2.3	2.2			
Sedative (non-medical use)	4.1	4.0	4.2	3.3	3.2	3.5	2.1	1.9	2.3	2.7	2.6	2.6	2.6
Tranquilizer (non-medical use)	8.3	8.5	8.2	6.2	5.8	6.3	3.5	3.2	3.6	3.9	4.0	4.2	4.7

\* "Binge" alcohol use in the National Survey on Drug Use and Health: Consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

\*\* "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: [54]