

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 12 AND OLDER, 2000-2010**

Estimated Past-Month Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	51.8	51.9	51.6	51.1	50.9	51.8	50.3	50.1	51.0	48.3	46.6
Binge Alcohol Use *	23.1	23.7	23.3	23.3	23.0	22.7	22.8	22.6	22.9	20.5	20.6
Heavy Alcohol Use **	6.7	6.8	6.9	6.9	6.9	6.6	6.9	6.8	6.7	5.7	5.6
Any Tobacco	27.4	27.7	28.4	28.6	29.6	29.4	29.2	29.8	30.4	29.5	29.3
Cigarettes	23.0	23.3	23.9	24.2	25.0	24.9	24.9	25.4	26.0	24.9	24.9
Smokeless Tobacco	3.5	3.4	3.5	3.2	3.3	3.2	3.0	3.3	3.3	3.2	3.4
Any Illicit Drug	8.9	8.7	8.0	8.0	8.3	8.1	7.9	8.2	8.3	7.1	6.3
Marijuana/Hashish	6.9	6.6	6.1	5.8	6.0	6.0	6.1	6.2	6.2	5.4	4.8
Inhalant	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.3
Hallucinogen	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4
LSD	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	NA
Cocaine (any form)	0.6	0.7	0.7	0.8	1.0	1.0	0.8	1.0	0.9	0.7	0.5
Crack Cocaine	0.1	0.2	0.1	0.2	0.3	0.3	0.2	0.3	0.2	0.2	0.1
Heroin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Psychotherapeutic Drugs (non-medical use)	2.7	2.8	2.5	2.8	2.9	2.7	2.5	2.7	2.7		
Pain Reliever (non-medical use)	2.0	2.1	1.9	2.1	2.1	1.9	1.8	2.0	1.9	1.6	1.2
OxyContin	0.2	0.2	0.2	0.1	0.1	0.1	0.1	NA	NA	NA	NA
Stimulant (non-medical use)	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.4
Methamphetamine	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.2
Sedative (non-medical use)	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1
Tranquilizer (non-medical use)	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.8	0.8	0.6	0.4

Estimated Lifetime Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	82.5	82.8	82.2	82.3	82.7	82.9	82.4	83.1	83.1	81.7	81.0
Any Tobacco	68.7	69.1	69.6	69.5	70.7	70.8	71.4	72.7	73.1	71.4	70.5
Cigarettes	64.2	64.6	65.1	65.3	66.3	66.6	67.3	68.7	69.1	67.2	66.5
Smokeless Tobacco	17.7	17.8	18.4	18.2	18.6	18.4	18.6	19.4	19.9	19.1	18.5
Any Illicit Drug	47.1	47.1	47.0	46.1	45.4	46.1	45.8	46.4	46.0	41.7	38.9
Marijuana/Hashish	42.0	41.7	41.0	40.6	39.8	40.1	40.2	40.6	40.4	36.9	34.2
Inhalant	8.6	9.0	8.9	9.1	9.3	9.4	9.5	9.7	9.7	8.1	7.5
Hallucinogen	14.8	14.8	14.4	13.8	14.3	13.9	14.3	14.5	14.6	12.5	11.7
LSD	9.2	9.4	9.4	9.1	9.5	9.2	9.7	10.3	10.4	9.0	8.8
PCP	2.5	2.5	2.7	2.5	2.7	2.7	2.8	3.0	3.2	2.7	2.6
Ecstasy	6.3	5.7	5.2	5.0	5.0	4.7	4.6	4.6	4.3	3.6	2.9
Cocaine (any form)	14.7	14.5	14.7	14.5	14.3	13.8	14.2	14.7	14.4	12.3	11.2
Crack Cocaine	3.6	3.3	3.4	3.5	3.5	3.3	3.3	3.3	3.6	2.8	2.4
Heroin	1.6	1.5	1.5	1.5	1.5	1.5	1.3	1.6	1.6	1.4	1.2
Psychotherapeutic Drugs (non-medical use)	20.4	20.6	20.8	20.3	20.7	20.4	20.4	20.6	20.4		
Pain Reliever (non-medical use)	13.7	13.9	14.0	13.3	13.6	13.4	13.2	13.1	12.6	9.8	8.6
OxyContin	2.4	2.3	1.9	1.8	1.7	1.4	1.3	1.2	0.8	NA	NA
Stimulant (non-medical use)	8.5	8.7	8.5	8.7	8.2	7.8	8.3	8.8	9.0	7.1	6.6
Methamphetamine	5.1	5.1	5.0	5.3	5.8	4.3	4.9	5.2	5.3	4.3	4.0
Sedative (non-medical use)	3.0	3.4	3.6	3.4	3.6	3.7	4.1	4.0	4.2	3.3	3.2
Tranquilizer (non-medical use)	8.7	8.6	8.6	8.2	8.7	8.7	8.3	8.5	8.2	6.2	5.8
Needle/Injection Drug Use	1.4	1.5	1.6	1.6	1.6	1.5	1.5	1.5	1.7	1.5	

NA Not included in survey for indicated year

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Data Source:

U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2k10ResultsTables/NSDUHTables2010R/HTM/TOC.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 12-17, 2000-2010**

Estimated Past-Month Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	13.6	14.7	14.6	15.9	16.6	16.5	17.6	17.7	17.6	17.3	16.4
Binge Alcohol Use *	7.8	8.8	8.8	9.7	10.3	9.9	11.1	10.6	10.7	10.6	10.4
Heavy Alcohol Use **	1.7	2.1	2.0	2.3	2.4	2.4	2.7	2.6	2.5	2.5	2.6
Any Tobacco	10.7	11.6	11.4	12.4	12.9	13.1	14.4	14.4	15.2	15.1	15.6
Cigarettes	8.3	8.9	9.1	9.8	10.4	10.8	11.9	12.2	13.0	13.0	13.4
Smokeless Tobacco	2.3	2.3	2.2	2.4	2.4	2.1	2.3	2.0	2.0	2.1	2.1
Any Illicit Drug	10.1	10.1	9.3	9.5	9.8	9.9	10.6	11.2	11.6	10.8	9.7
Marijuana/Hashish	7.4	7.3	6.7	6.7	6.7	6.8	7.6	7.9	8.2	8.0	7.2
Inhalant	1.1	1.0	1.1	1.2	1.3	1.2	1.2	1.3	1.2	1.0	1.0
Hallucinogen	0.9	0.9	1.0	0.7	0.7	0.8	0.8	1.0	1.0	1.2	1.2
LSD	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.4	0.5
PCP	0.0	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Ecstasy	0.5	0.5	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.6	NA
Cocaine (any form)	0.2	0.3	0.4	0.4	0.4	0.6	0.5	0.6	0.6	0.4	0.6
Crack Cocaine	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Heroin	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.1
Psychotherapeutic Drugs (non-medical use)	3.0	3.1	2.9	3.3	3.3	3.3	3.6	4.0	4.0		
Pain Reliever (non-medical use)	2.5	2.7	2.3	2.7	2.7	2.7	3.0	3.2	3.2	2.6	2.3
OxyContin	0.2	0.3	0.2	0.2	0.1	0.1	0.3	NA	NA	NA	NA
Stimulant (non-medical use)	0.4	0.5	0.5	0.5	0.7	0.7	0.7	0.9	0.8	0.7	0.8
Methamphetamine	0.1	0.1	0.1	0.1	0.2	0.3	0.2	0.3	0.3	0.2	0.3
Sedative (non-medical use)	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.2	0.1	0.2
Tranquilizer (non-medical use)	0.5	0.6	0.6	0.7	0.5	0.6	0.6	0.9	0.8	0.5	0.5

Estimated Lifetime Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	35.2	38.1	38.3	39.4	40.4	40.6	42.0	42.9	43.4	42.9	41.7
Any Tobacco	24.9	26.9	27.2	27.8	29.7	30.7	32.7	34.5	36.8	37.0	38.0
Cigarettes	20.3	22.2	22.9	23.7	25.8	26.7	29.2	31.0	33.3	33.6	34.6
Smokeless Tobacco	7.2	7.6	7.2	7.7	7.1	7.1	7.1	7.6	8.0	8.3	8.6
Any Illicit Drug	25.7	26.8	26.2	26.2	27.6	27.7	30.0	30.5	30.9	28.4	26.9
Marijuana/Hashish	17.0	17.0	16.5	16.2	17.3	17.4	19.0	19.6	20.6	19.7	18.3
Inhalant	8.2	9.2	9.3	9.6	10.1	10.5	11.0	10.7	10.5	8.6	8.9
Hallucinogen	4.1	4.1	3.9	3.5	3.9	3.9	4.6	5.0	5.7	5.7	5.8
LSD	0.9	1.0	1.1	0.8	0.9	1.1	1.2	1.6	2.7	3.1	3.6
PCP	0.3	0.4	0.4	0.4	0.5	0.7	0.7	0.8	0.9	1.0	1.1
Ecstasy	2.5	2.3	2.1	1.8	1.9	1.6	2.1	2.4	3.3	3.2	2.6
Cocaine (any form)	1.5	1.6	1.9	2.1	2.2	2.3	2.4	2.6	2.7	2.3	2.4
Crack Cocaine	0.2	0.2	0.3	0.4	0.5	0.4	0.5	0.6	0.7	0.6	0.6
Heroin	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.3	0.4	0.3	0.4
Psychotherapeutic Drugs (non-medical use)	10.6	11.4	11.1	11.7	12.5	11.9	13.5	13.5	13.7		
Pain Reliever (non-medical use)	9.2	9.7	9.4	9.7	10.4	9.9	11.4	11.2	11.2	9.4	8.4
OxyContin	1.5	1.5	1.4	1.4	1.3	1.1	1.2	1.0	0.9	NA	NA
Stimulant (non-medical use)	2.2	2.4	2.7	2.9	3.7	3.4	3.6	4.2	4.5	3.7	4.0
Methamphetamine	0.6	0.8	0.8	0.9	1.3	1.3	1.4	1.5	1.7	1.4	1.3
Sedative (non-medical use)	0.7	0.9	0.8	0.8	0.9	0.9	1.0	1.0	1.0	0.7	0.8
Tranquilizer (non-medical use)	3.0	2.8	2.9	3.0	2.9	3.0	3.2	3.5	3.4	2.6	2.5
Needle/Injection Drug Use	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.3	0.3		

NA Not included in survey for indicated year

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Data Source:

U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2k10ResultsTables/NSDUHTables2010R/HTM/TOC.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 18-25, 2000-2010**

Estimated Past-Month Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	61.5	61.8	61.2	61.2	61.9	60.9	60.5	61.4	60.5	58.8	56.8
Binge Alcohol Use *	40.6	41.7	41.0	41.8	42.2	41.9	41.2	41.6	40.9	38.7	37.8
Heavy Alcohol Use **	13.6	13.7	14.5	14.7	15.6	15.3	15.1	15.1	14.9	13.6	12.8
Any Tobacco	40.8	41.6	41.4	41.8	43.9	44.3	44.6	44.8	45.3	43.9	42.9
Cigarettes	34.2	35.8	35.7	36.2	38.4	39.0	39.5	40.2	40.8	39.1	38.3
Smokeless Tobacco	6.4	6.1	5.4	5.3	5.2	5.1	4.9	4.7	4.8	5.4	5.0
Any Illicit Drug	21.5	21.2	19.6	19.7	19.8	20.1	19.4	20.3	20.2	18.8	15.9
Marijuana/Hashish	18.5	18.1	16.5	16.4	16.3	16.6	16.1	17.0	17.3	16.0	13.6
Inhalant	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.4	0.5	0.6	0.6
Hallucinogen	2.0	1.8	1.7	1.5	1.7	1.5	1.5	1.7	1.9	2.7	1.8
LSD	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.1	0.6	0.8
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.1
Ecstasy	1.2	1.1	0.9	0.7	1.0	0.8	0.7	0.7	1.1	1.7	NA
Cocaine (any form)	1.5	1.4	1.5	1.7	2.2	2.6	2.1	2.2	2.0	1.9	1.4
Crack Cocaine	0.2	0.1	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.1
Heroin	0.3	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.2	0.1
Psychotherapeutic Drugs (non-medical use)	5.9	6.3	5.9	6.0	6.5	6.3	6.1	6.1	5.5		
Pain Reliever (non-medical use)	4.5	4.8	4.6	4.6	4.9	4.7	4.7	4.7	4.1	3.6	2.7
OxyContin	0.7	0.5	0.4	0.5	0.4	0.4	0.4	NA	NA	NA	NA
Stimulant (non-medical use)	1.2	1.3	1.1	1.1	1.4	1.4	1.5	1.3	1.3	1.3	0.8
Methamphetamine	0.2	0.2	0.2	0.4	0.6	0.7	0.7	0.6	0.6	0.7	0.3
Sedative (non-medical use)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
Tranquilizer (non-medical use)	1.7	1.8	1.7	1.7	2.0	1.9	1.8	1.7	1.6	1.3	1.0

Estimated Lifetime Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	85.7	85.8	85.6	85.2	86.5	85.7	86.2	87.1	86.7	85.0	84.0
Any Tobacco	68.6	69.6	70.2	70.3	71.8	72.3	73.5	74.8	75.5	73.6	72.1
Cigarettes	62.1	63.7	64.2	64.7	66.6	67.3	68.7	70.2	71.2	69.0	67.3
Smokeless Tobacco	20.9	21.0	20.3	20.1	20.2	20.8	21.4	22.0	23.7	23.7	23.6
Any Illicit Drug	57.0	58.1	56.6	57.4	59.0	59.2	59.2	60.5	59.8	55.6	51.2
Marijuana/Hashish	51.1	52.2	50.4	50.8	52.4	52.4	52.8	53.9	53.8	50.0	45.7
Inhalant	9.9	10.7	10.4	11.3	12.5	13.3	14.0	14.9	15.7	13.4	12.8
Hallucinogen	18.4	19.0	17.7	19.2	20.2	21.0	21.3	23.3	24.2	22.1	19.3
LSD	6.4	6.8	6.5	7.3	8.9	10.5	12.1	14.0	15.9	15.3	14.0
PCP	1.3	1.4	1.6	1.6	1.9	2.2	2.3	3.0	2.7	2.6	2.3
Ecstasy	12.4	12.4	12.1	12.8	13.4	13.7	13.8	14.8	15.1	13.1	9.7
Cocaine (any form)	13.3	14.9	14.4	15.0	15.7	15.1	15.2	15.0	15.4	13.0	10.9
Crack Cocaine	2.6	2.9	2.7	3.0	3.6	3.5	3.5	3.8	3.8	3.4	2.8
Heroin	1.8	1.7	1.4	1.5	1.6	1.5	1.6	1.6	1.6	1.6	1.4
Psychotherapeutic Drugs (non-medical use)	28.7	29.2	29.2	29.9	30.7	30.8	29.7	29.5	28.3		
Pain Reliever (non-medical use)	23.9	24.5	24.4	24.8	25.5	25.5	24.3	23.7	22.1	18.2	14.6
OxyContin	6.2	6.2	5.3	5.2	5.1	5.0	4.3	3.6	2.6	NA	NA
Stimulant (non-medical use)	10.3	10.6	10.5	10.9	12.0	12.3	12.1	12.2	12.2	9.5	7.6
Methamphetamine	4.0	4.5	4.7	5.6	6.4	6.7	7.0	6.9	7.4	5.1	4.1
Sedative (non-medical use)	1.4	1.6	1.5	1.8	1.9	2.0	1.9	1.8	2.1	1.9	1.6
Tranquilizer (non-medical use)	13.1	12.8	12.2	12.7	13.1	13.4	12.2	12.3	11.2	8.9	7.4
Needle/Injection Drug Use	1.1	1.3	0.9	1.3	1.2	1.4	1.3	1.3	1.2		

NA Not included in survey for indicated year

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Data Source:

U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2k10ResultsTables/NSDUHTables2010R/HTM/TOC.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 26 AND OLDER, 2000-2010**

Estimated Past-Month Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	54.9	54.9	54.8	54.1	53.7	55.1	53.0	52.5	53.9	50.8	49.0
Binge Alcohol Use *	21.9	22.4	22.1	21.9	21.4	21.0	21.1	21.0	21.4	18.8	19.1
Heavy Alcohol Use **	6.1	6.2	6.3	6.1	6.0	5.6	6.1	5.9	5.9	4.8	4.8
Any Tobacco	27.2	27.3	28.3	28.5	29.4	29.0	28.5	29.3	29.9	29.0	28.9
Cigarettes	22.8	23.0	23.8	24.1	24.7	24.3	24.1	24.7	25.2	24.2	24.2
Smokeless Tobacco	3.1	3.1	3.3	3.0	3.2	3.0	2.7	3.2	3.2	3.0	3.3
Any Illicit Drug	6.6	6.3	5.9	5.8	6.1	5.8	5.5	5.6	5.8	4.5	4.2
Marijuana/Hashish	4.8	4.6	4.2	3.9	4.2	4.1	4.1	4.0	4.0	3.2	3.0
Inhalant	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogen	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.1
LSD	NE	0.0	NE	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCP	0.0	0.0	NE	0.0	0.0	0.0	0.0	NE	0.0	NE	0.0
Ecstasy	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	NA
Cocaine (any form)	0.5	0.6	0.7	0.7	0.8	0.8	0.7	0.8	0.7	0.6	0.4
Crack Cocaine	0.2	0.2	0.2	0.3	0.3	0.3	0.2	0.3	0.3	0.2	0.1
Heroin	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.0	0.1
Psychotherapeutic Drugs (non-medical use)	2.2	2.1	1.9	2.2	2.2	1.9	1.8	2.0	2.0		
Pain Reliever (non-medical use)	1.5	1.6	1.4	1.6	1.5	1.3	1.2	1.3	1.3	1.1	0.9
OxyContin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	NA	NA	NA	NA
Stimulant (non-medical use)	0.3	0.4	0.2	0.3	0.4	0.3	0.4	0.4	0.4	0.3	0.2
Methamphetamine	0.1	0.2	0.1	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.1
Sedative (non-medical use)	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1
Tranquilizer (non-medical use)	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.6	0.6	0.5	0.4

Estimated Lifetime Substance Use (%)

	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001	2000
Alcohol	87.9	88.0	87.3	87.5	87.7	88.2	87.3	88.0	88.0	86.5	85.8
Any Tobacco	74.2	74.4	75.0	75.0	76.0	76.0	76.4	77.6	77.7	75.7	74.7
Cigarettes	70.0	70.1	70.7	70.9	71.7	71.9	72.3	73.6	73.7	71.5	70.7
Smokeless Tobacco	18.5	18.5	19.5	19.3	19.9	19.5	19.6	20.6	20.9	19.8	19.1
Any Illicit Drug	48.1	47.8	48.0	46.8	45.5	46.3	45.6	46.1	45.7	41.2	38.5
Marijuana/Hashish	43.4	42.7	42.6	42.0	40.6	41.1	41.0	41.2	40.8	37.0	34.4
Inhalant	8.3	8.6	8.6	8.6	8.6	8.5	8.5	8.6	8.6	7.1	6.4
Hallucinogen	15.5	15.4	15.2	14.2	14.7	14.0	14.4	14.2	14.1	11.9	11.2
LSD	10.7	10.9	11.0	10.6	10.8	10.1	10.5	10.8	10.5	8.7	8.6
PCP	2.9	2.9	3.1	2.9	3.1	3.1	3.2	3.3	3.5	2.9	2.9
Ecstasy	5.7	4.9	4.4	4.1	3.9	3.6	3.4	3.1	2.6	2.0	1.8
Cocaine (any form)	16.6	16.1	16.4	16.0	15.8	15.2	15.6	16.3	15.9	13.6	12.4
Crack Cocaine	4.2	3.8	3.9	3.9	3.9	3.6	3.6	3.6	3.9	2.9	2.5
Heroin	1.8	1.6	1.7	1.7	1.7	1.6	1.4	1.7	1.7	1.5	1.3
Psychotherapeutic Drugs (non-medical use)	20.1	20.2	20.6	19.8	20.1	19.7	19.8	20.0	19.9		
Pain Reliever (non-medical use)	12.6	12.7	12.7	11.8	12.0	11.8	11.5	11.5	11.1	8.4	7.6
OxyContin	1.9	1.7	1.4	1.2	1.1	0.9	0.8	0.8	0.5	NA	NA
Stimulant (non-medical use)	9.0	9.2	8.9	9.1	9.4	8.7	9.6	10.0	10.4	7.1	6.7
Methamphetamine	5.9	5.8	5.7	5.8	6.3	5.5	6.5	6.9	7.1	4.5	4.3
Sedative (non-medical use)	3.6	4.1	4.3	4.0	4.3	4.4	4.9	4.8	5.1	3.9	3.8
Tranquilizer (non-medical use)	8.7	8.7	8.7	8.1	8.7	8.6	8.3	8.5	8.3	6.2	6.0
Needle/Injection Drug Use	1.6	1.7	2.0	1.9	1.8	1.7	1.6	1.8	2.0		

NA Not included in survey for indicated year

NE No estimate reported due to low precision

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.


Data Source:

U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2k10ResultsTables/NSDUHTables2010R/HTM/TOC.htm>


ALCOHOL — PAST-MONTH USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12 and Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL — PAST-MONTH BINGE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL — PERCEIVED RISK OF BINGE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


TOBACCO — PAST-MONTH CIGARETTE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


TOBACCO — PERCEIVED RISK OF CIGARETTE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12 & Older, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12-17, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 18-25, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 26 & Older, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAST-MONTH ILLICIT DRUG USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PAST-MONTH USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PAST-YEAR USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 12 & Older, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 12-17, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 18-25, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 26 & Older, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PERCEIVED RISK OF USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


COCAINE — PAST-YEAR USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAIN RELIEVERS: PAST-YEAR NON-MEDICAL USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12 & Older, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12-17, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 18-25, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 26 & Older, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL AND ILLICIT DRUGS — PAST-YEAR DEPENDENCE OR ABUSE: U.S. AND MISSOURI AGE GROUPS

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%), U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

ANY MENTAL ILLNESS IN PAST YEAR: U.S. AND MISSOURI COMPARISON BY AGE GROUP


Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 18 & Older, 2009-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 18-25, 2009-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 26 and Older, 2009-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

SERIOUS MENTAL ILLNESS IN PAST YEAR: U.S. AND MISSOURI COMPARISON BY AGE GROUP


Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 18 & Older, 2009-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 18-25, 2009-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 26 and Older, 2009-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


DEPRESSION — MAJOR PAST-YEAR EPISODE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 12-17, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18 and Older, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18-25, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 26 and Older, 2005-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.


PAST-MONTH ALCOHOL USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12 and Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


PAST-MONTH BINGE ALCOHOL USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


PERCEIVED RISK OF BINGE DRINKING (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk Binge Drinking Weekly (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Perceive Great Risk Binge Drinking Weekly (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Perceive Great Risk Binge Drinking Weekly (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


Perceive Great Risk Binge Drinking Weekly (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


ALCOHOL USE, BINGE USE, AND PERCEIVED RISK (%): MISSOURI AGE GROUP COMPARISONS

Estimated Past-Month Alcohol Use (%): Missouri Age Group Comparisons, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.

Estimated Past-Month Binge Drinking (%): Missouri Age Group Comparisons, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


Perceptions of Great Risk of Binge Drinking (%): Missouri Age Group Comparisons, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. Estimates from the National Survey on Drug Use and Health.


ALCOHOL USE, BINGE USE, AND PERCEIVED RISK: 3-YEAR AVERAGES FOR MISSOURI AND REGIONS

Past-Month Alcohol Use (%): Missouri and ADA/CPS Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Past-Month Binge Drinking (%): Missouri and ADA/CPS Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Perceptions of Great Risk of Binge Drinking (%): Missouri and ADA/CPS Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


PAST 30-DAY ALCOHOL USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

2011 YRBS Missouri data not available on publication date.


PAST 30-DAY BINGE ALCOHOL USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

2011 YRBS Missouri data not available on publication date.


PAST 30-DAY, BINGE, AND EARLY ALCOHOL USE: MISSOURI GRADE 9-12 COMPARISONS, 1995-2009

Past 30-Day Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey


Alcohol Use Before Age 13 (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey


ALCOHOL USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Alcohol (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Alcohol (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 2-Week (2004-2008) and Past 30-Day (2010-2012) Binge Alcohol Use (%): Missouri Students in Grades 6-12



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


PAST-MONTH ALCOHOL USE (%): MISSOURI AGE GROUP AND GRADE LEVEL COMPARISONS

Estimated Past-Month Alcohol Use (%): Missouri Age Group Comparisons, 2001-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey


Reported Past 30-Day Alcohol Use (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


PAST YEAR, PAST 30-DAY, AND BINGE ALCOHOL USE: MISSOURI COLLEGE STUDENTS, 2007-2012

Past-Year Alcohol Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.

Past 30-Day Alcohol Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.

Past 2-Week Binge Alcohol Use in One Sitting (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.


Past 2-Week Binge Alcohol Use in 2-Hour Period (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.


PAST-MONTH CIGARETTE USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PERCEIVED RISK OF CIGARETTE USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2003-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Perceive Great Risk Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2003-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Perceive Great Risk Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2003-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Perceive Great Risk Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2003-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PAST-MONTH TOBACCO USE AND PERCEIVED RISK: 3-YEAR AVERAGES FOR MISSOURI AND REGIONS

Estimated Past-Month Tobacco Use (%): Missouri and ADA/CPS Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Month Cigarette Use (%): Missouri and ADA/CPS Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Perceive Great Risk in Smoking One or More Packs of Cigarettes Per Day (%): Missouri and ADA/CPS Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


PAST 30-DAY CIGARETTE USE (%): U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

2011 YRBS Missouri data not available on publication date.


TOBACCO USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Cigarettes (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Cigarettes (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Lifetime Use of Chewing Tobacco (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Chewing Tobacco (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


PAST-MONTH CIGARETTE USE (%): MISSOURI AGE GROUP AND GRADE LEVEL COMPARISONS

Estimated Past-Month Cigarette Use (%): Missouri Age Group Comparisons, 2-Year Averages, 2001-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Cigarette Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey


Reported Past 30-Day Cigarette Use (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


PAST-MONTH ILLICIT DRUG USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Use of any Illicit Drug (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Use of any Illicit Drug (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Use of any Illicit Drug (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Month Use of any Illicit Drug (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PAST-MONTH MARIJUANA USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PERCEIVED RISK OF MARIJUANA USE (%): U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12 & Older, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Perceive Great Risk Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12-17, 2-Year Averages, 2000-2009




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Perceive Great Risk Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 18-25, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Perceive Great Risk Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 26 & Older, 2-Year Averages, 2000-2009



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PAST-YEAR ALCOHOL OR ILLICIT DRUG DEPENDENCE OR ABUSE: MISSOURI AGE GROUPS

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12-17, 2001-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 18-25, 2001-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PAST-MONTH, FIRST USE, AND PERCEIVED RISK OF MARIJUANA: 3-YEAR AVERAGES FOR MISSOURI AND REGIONS

Estimated Past-Month Marijuana Use (%): Missouri and ADA Regions for Combined Years




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Average Annual Rate of First Use of Marijuana (%): Missouri and ADA Regions for Combined Years



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Perception of Great Risk of Smoking Marijuana Monthly (%): Missouri and ADA Regions for Combined Years



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


MARIJUANA USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Marijuana (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Marijuana (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


PAST-MONTH MARIJUANA USE (%): MISSOURI AGE GROUP AND GRADE LEVEL COMPARISONS

Estimated Past-Month Marijuana Use (%): Missouri Age Group Comparisons, 2001-2008




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Marijuana Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey


Reported Past 30-Day Marijuana Use (%): Missouri Students in Grades 6-12, 2004-2010



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


COCAINE AND HEROIN USE: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Cocaine (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Cocaine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


Lifetime Use of Heroin (%): Missouri Students in Grades 6-12, 2010-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


PAST-YEAR NON-MEDICAL USE OF PAIN RELIEVERS: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12 & Older, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12-17, 2003-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 18-25, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 26 & Older, 2003-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


METHAMPHETAMINE USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Speed, Amphetamine or Methamphetamine (%): Missouri Students in Grades 6-12, 2004-2008




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.

Lifetime Use of Methamphetamine (%): Missouri Students in Grades 6-12, 2010-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Past 30-Day Use of Speed, Amphetamine or Methamphetamine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


INHALANT USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Inhalants (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Inhalants (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


HALLUCINOGEN USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Hallucinogens such as LSD, PCP, Mescaline, Peyote (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of LSD or other Psychedelics (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


CLUB DRUG USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Club Drugs such as Ecstasy, GHB, Rohypnol, Ketamine (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Club Drugs such as Ecstasy, GHB, Rohypnol, Ketamine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


PHARMACEUTICAL DRUG USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Non-Prescribed Pharmaceutical Drugs (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Past 30-Day Use of Non-Prescribed Pharmaceutical Drugs (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Lifetime Use of Non-Prescribed Steroids (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


OVER-THE-COUNTER AND SYNTHETIC DRUG USE: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Over-the-Counter Medicines for Non-medical Reason (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Past 30-Day Use of Over-the-Counter Meds for Non-medical Reason (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Lifetime Use of Synthetic Drugs such as K2, Bath Salts, Spice (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Past 30-Day Use of Synthetic Drugs such as K2, Bath Salts, Spice (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

LIFETIME INJECTION DRUG USE: MISSOURI STUDENTS IN GRADES 6-12


Lifetime Injection Drug Use (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. 2004-2008 Missouri Student Survey respondents. Data are unweighted.


PAST-YEAR ILLICIT DRUG USE: MISSOURI COLLEGE STUDENTS, 2007-2012

Past-Year Marijuana Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.

Past-Year Cocaine Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents. Item not included on 2012 survey.

Past-Year Amphetamine Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents. Item not included on 2012 survey.

Past-Year Inhalant Use (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents. Item not included on 2012 survey.


Past-Year Club Drug Use (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents. Item not included on 2012 survey.


PAST-YEAR ALCOHOL OR ILLICIT DRUG DEPENDENCE OR ABUSE: U.S. AND MISSOURI AGE GROUPS

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%), U.S. and Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 12-17, 2001-2010




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 18-25, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAST-YEAR ALCOHOL OR ILLICIT DRUG DEPENDENCE OR ABUSE: MISSOURI AGE GROUPS

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12 & Older, 2001-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12-17, 2001-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 18-25, 2001-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 26 & Older, 2001-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

PAST-YEAR SUBSTANCE DEPENDENCE OR ABUSE: MISSOURI AND REGIONS


Estimated Past-Year Alcohol Dependence or Abuse (%): Missouri and ADA/CPS Regions for Combined Years



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are annual averages based on 3-year samples


Estimated Past-Year Illicit Drug Dependence or Abuse (%): Missouri and ADA/CPS Regions for Combined Years



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are annual averages based on 3-year samples

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): Missouri and ADA/CPS Regions for Combined Years




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are annual averages based on 3-year samples


PAST-YEAR UNMET NEED FOR SUBSTANCE ABUSE TREATMENT: MISSOURI AGE GROUPS

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 12 & Older, 2002-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 12-17, 2002-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 18-25, 2002-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 26 & Older, 2002-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


PAST-YEAR UNMET NEED FOR SUBSTANCE ABUSE TREATMENT: 3-YEAR AVERAGES FOR MISSOURI AND REGIONS

Past-Year Percentage Needing but Not Receiving Treatment for Alcohol Use: Missouri & ADA/CPS Regions, 3-Year Spans



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.
Estimates are annual averages based on 3-year samples.


Past-Year Percentage Needing but Not Receiving Treatment for Illicit Drug Use: Missouri & ADA/CPS Regions, 3-Year Spans



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.
Estimates are annual averages based on 3-year samples.


PAST-YEAR MAJOR DEPRESSIVE EPISODE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 12-17, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18 and Older, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 not comparable to prior years.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18-25, 2005-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 not comparable to prior years.

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 26 and Older, 2005-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 not comparable to prior years.


MAJOR DEPRESSIVE EPISODE IN PAST YEAR: MISSOURI AND ADA/CPS REGIONS

Estimated Past-Year Major Depressive Episode (MDE): Ages 12-17, Missouri and ADA/CPS Regions, 3-Year Span



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Year Major Depressive Episode (MDE): Ages 18 and Older, Missouri and ADA/CPS Regions, 3-Year Spans



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.