

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 12 AND OLDER, 2002-2012**

Estimated Past-Month Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	52.1	51.8	51.8	51.9	51.6	51.2	51.0	51.8	50.3	50.1	51.0
Binge Alcohol Use *	23.0	22.6	23.1	23.7	23.4	23.3	23.0	22.7	22.8	22.6	22.9
Heavy Alcohol Use **	6.5	6.2	6.7	6.8	7.0	6.9	6.9	6.6	6.9	6.8	6.7
Any Tobacco	26.7	26.5	27.5	27.7	28.4	28.7	29.6	29.4	29.2	29.8	30.4
Cigarettes	22.1	22.1	23.0	23.3	24.0	24.3	25.0	24.9	24.9	25.4	26.0
Smokeless Tobacco	3.5	3.2	3.5	3.4	3.5	3.3	3.3	3.2	3.0	3.3	3.3
Any Illicit Drug	9.2	8.7	8.9	8.7	8.1	8.0	8.3	8.1	7.9	8.2	8.3
Marijuana	7.3	7.0	6.9	6.7	6.1	5.8	6.0	6.0	6.1	6.2	6.2
Inhalant	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.2	0.3
Hallucinogen	0.4	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.5
LSD	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.3
Cocaine (any form)	0.6	0.5	0.6	0.7	0.7	0.8	1.0	1.0	0.8	1.0	0.9
Crack Cocaine	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.3	0.2	0.3	0.2
Heroin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Psychotherapeutics (Rx-type, non-medical use)	2.6	2.4	2.7	2.8	2.5	2.8	2.9	2.7	2.5	2.7	2.7
Pain Reliever	1.9	1.7	2.0	2.1	1.9	2.1	2.1	1.9	1.8	2.0	1.9
OxyContin®	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1		
Stimulant	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.5	0.5
Methamphetamine	0.2	0.2	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.3	0.3
Sedative	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2
Tranquilizer	0.8	0.7	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.8	0.8

Estimated Lifetime Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	82.3	82.2	82.5	82.8	82.2	82.3	82.7	82.9	82.4	83.1	83.1
Any Tobacco	66.8	67.5	68.7	69.1	69.6	69.5	70.7	70.8	71.4	72.7	73.1
Cigarettes	61.9	62.8	64.2	64.6	65.1	65.3	66.3	66.6	67.3	68.7	69.1
Smokeless Tobacco	17.7	18.0	17.7	17.8	18.4	18.2	18.6	18.4	18.6	19.4	19.9
Any Illicit Drug	48.0	47.0	47.3	47.4	47.2	46.2	45.4	46.1	45.8	46.4	46.0
Marijuana	42.8	41.9	42.0	41.7	41.2	40.6	39.7	40.1	40.2	40.6	40.4
Inhalant	8.1	8.0	8.6	9.0	9.0	9.1	9.3	9.4	9.5	9.7	9.7
Hallucinogen	14.6	14.1	14.8	14.9	14.4	13.8	14.3	13.9	14.3	14.5	14.6
LSD	9.1	8.9	9.2	9.4	9.4	9.2	9.5	9.2	9.7	10.3	10.4
PCP	2.5	2.4	2.5	2.5	2.7	2.5	2.7	2.7	2.8	3.0	3.2
Ecstasy	6.2	5.7	6.3	5.7	5.2	5.0	5.0	4.7	4.6	4.6	4.3
Cocaine (any form)	14.5	14.3	14.7	14.6	14.8	14.5	14.3	13.8	14.2	14.7	14.4
Crack Cocaine	3.5	3.2	3.6	3.3	3.4	3.5	3.5	3.3	3.3	3.3	3.6
Heroin	1.8	1.6	1.6	1.5	1.5	1.5	1.5	1.5	1.3	1.6	1.6
Psychotherapeutics (Rx-type, non-medical use)	20.9	19.9	20.4	20.6	20.9	20.4	20.7	20.4	20.4	20.6	20.4
Pain Reliever	14.2	13.3	13.8	14.0	14.0	13.4	13.6	13.4	13.2	13.1	12.6
OxyContin®	2.5	2.3	2.4	2.3	1.9	1.8	1.7	1.4	1.3	1.2	0.8
Stimulant	8.3	7.9	8.6	8.7	8.5	8.7	8.2	7.8	8.3	8.8	9.0
Methamphetamine	4.7	4.6	5.1	5.1	5.1	5.3	5.8	4.3	4.9	5.2	5.3
Sedative	3.1	2.9	3.0	3.4	3.6	3.4	3.6	3.7	4.1	4.0	4.2
Tranquilizer	9.1	8.4	8.7	8.7	8.6	8.2	8.7	8.7	8.3	8.5	8.2
Needle/Injection Drug Use	1.6	1.4	1.4	1.5	1.6	1.6	1.6	1.5	1.5	1.5	1.7

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Some estimates from 2006-2010 differ slightly from those published in previous Status Reports due to recent SAMHSA updates.

Data Source:

U.S. Department of Health and Human Services, SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DetTabs/NSDUH-DetTabsTOC2012.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 12-17, 2002-2012**

Estimated Past-Month Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	12.9	13.3	13.6	14.8	14.7	16.0	16.7	16.5	17.6	17.7	17.6
Binge Alcohol Use *	7.2	7.4	7.9	8.9	8.9	9.7	10.3	9.9	11.1	10.6	10.7
Heavy Alcohol Use **	1.3	1.5	1.7	2.1	2.0	2.3	2.4	2.4	2.7	2.6	2.5
Any Tobacco	8.6	10.0	10.7	11.8	11.5	12.4	12.9	13.1	14.4	14.4	15.2
Cigarettes	6.6	7.8	8.4	9.0	9.2	9.9	10.4	10.8	11.9	12.2	13.0
Smokeless Tobacco	2.1	2.1	2.3	2.4	2.2	2.5	2.4	2.1	2.3	2.0	2.0
Any Illicit Drug	9.5	10.1	10.1	10.1	9.3	9.5	9.8	9.9	10.6	11.2	11.6
Marijuana	7.2	7.9	7.4	7.4	6.7	6.7	6.7	6.8	7.6	7.9	8.2
Inhalant	0.8	0.9	1.1	1.0	1.1	1.2	1.3	1.2	1.2	1.3	1.2
Hallucinogen	0.6	0.9	0.9	0.9	1.0	0.7	0.7	0.8	0.8	1.0	1.0
LSD	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.2
PCP	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1
Ecstasy	0.3	0.4	0.5	0.5	0.4	0.3	0.3	0.3	0.3	0.4	0.5
Cocaine (any form)	0.1	0.3	0.2	0.3	0.4	0.4	0.4	0.6	0.5	0.6	0.6
Crack Cocaine	***	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Heroin	***	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.0
Psychotherapeutics (Rx-type, non-medical use)	2.8	2.8	3.0	3.1	2.9	3.3	3.3	3.3	3.6	4.0	4.0
Pain Reliever	2.2	2.3	2.5	2.7	2.3	2.7	2.7	2.7	3.0	3.2	3.2
OxyContin®	0.2	0.2	0.2	0.3	0.2	0.2	0.1	0.1	0.3		
Stimulant	0.5	0.4	0.4	0.5	0.5	0.5	0.7	0.7	0.7	0.9	0.8
Methamphetamine	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.3	0.2	0.3	0.3
Sedative	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.2
Tranquilizer	0.6	0.6	0.5	0.6	0.6	0.7	0.5	0.6	0.6	0.9	0.8

Estimated Lifetime Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	32.4	34.5	35.2	38.1	38.3	39.4	40.4	40.6	42.0	42.9	43.4
Any Tobacco	21.5	23.2	24.9	26.9	27.2	27.8	29.7	30.7	32.7	34.5	36.8
Cigarettes	17.4	19.1	20.3	22.2	22.9	23.7	25.8	26.7	29.2	31.0	33.3
Smokeless Tobacco	6.4	6.8	7.2	7.6	7.2	7.7	7.1	7.1	7.1	7.6	8.0
Any Illicit Drug	24.2	25.5	25.8	26.9	26.4	26.3	27.6	27.7	30.0	30.5	30.9
Marijuana	17.0	17.5	17.1	17.1	16.6	16.2	17.3	17.4	19.0	19.6	20.6
Inhalant	6.5	7.5	8.3	9.3	9.3	9.6	10.1	10.5	11.0	10.7	10.5
Hallucinogen	3.3	3.7	4.1	4.1	3.9	3.6	3.9	3.9	4.6	5.0	5.7
LSD	1.0	0.9	0.9	1.0	1.1	0.8	0.9	1.1	1.2	1.6	2.7
PCP	0.4	0.3	0.3	0.4	0.4	0.4	0.5	0.7	0.7	0.8	0.9
Ecstasy	2.0	2.4	2.5	2.3	2.1	1.8	1.9	1.6	2.1	2.4	3.3
Cocaine (any form)	1.1	1.3	1.5	1.6	1.9	2.2	2.2	2.3	2.4	2.6	2.7
Crack Cocaine	0.1	0.3	0.2	0.2	0.3	0.4	0.5	0.4	0.5	0.6	0.7
Heroin	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.3	0.4
Psychotherapeutics (Rx-type, non-medical use)	10.0	10.1	10.6	11.4	11.2	11.7	12.5	11.9	13.5	13.5	13.7
Pain Reliever	8.3	8.6	9.2	9.7	9.5	9.7	10.4	9.9	11.4	11.2	11.2
OxyContin®	1.3	1.5	1.5	1.5	1.4	1.4	1.4	1.1	1.2	1.0	0.9
Stimulant	2.0	2.0	2.2	2.4	2.7	2.9	3.7	3.4	3.6	4.2	4.5
Methamphetamine	0.6	0.7	0.6	0.8	0.8	0.9	1.3	1.3	1.4	1.5	1.7
Sedative	0.7	0.6	0.7	0.9	0.8	0.8	0.9	0.9	1.0	1.0	1.0
Tranquilizer	2.8	2.8	3.0	2.8	2.9	3.0	2.9	3.0	3.2	3.5	3.4
Needle/Injection Drug Use	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.3	0.3

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

*** Low precision, no estimate available

Some estimates from 2006-2010 differ slightly from those published in previous Status Reports due to recent SAMHSA updates.

Data Source:

U.S. Department of Health and Human Services, SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DetTabs/NSDUH-DetTabsTOC2012.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 18-25, 2002-2012**

Estimated Past-Month Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	60.2	60.7	61.4	61.8	61.1	61.3	62.0	60.9	60.5	61.4	60.5
Binge Alcohol Use *	39.5	39.8	40.5	41.8	41.2	41.9	42.3	41.9	41.2	41.6	40.9
Heavy Alcohol Use **	12.7	12.1	13.5	13.8	14.6	14.8	15.6	15.3	15.1	15.1	14.9
Any Tobacco	38.1	39.5	40.9	41.6	41.4	41.9	44.0	44.3	44.6	44.8	45.3
Cigarettes	31.8	33.5	34.3	35.8	35.7	36.2	38.5	39.0	39.5	40.2	40.8
Smokeless Tobacco	5.5	5.4	6.4	6.1	5.3	5.2	5.2	5.1	4.9	4.7	4.8
Any Illicit Drug	21.3	21.4	21.6	21.4	19.7	19.8	19.8	20.1	19.4	20.3	20.2
Marijuana	18.7	19.0	18.5	18.2	16.6	16.5	16.3	16.6	16.1	17.0	17.3
Inhalant	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.4	0.5
Hallucinogen	1.7	1.6	2.0	1.8	1.7	1.5	1.7	1.5	1.5	1.7	1.9
LSD	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
Ecstasy	1.0	0.9	1.2	1.1	0.9	0.7	1.0	0.8	0.7	0.7	1.1
Cocaine (any form)	1.1	1.4	1.5	1.4	1.6	1.7	2.2	2.6	2.1	2.2	2.0
Crack Cocaine	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.3	0.3	0.2	0.2
Heroin	0.4	0.3	0.3	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1
Psychotherapeutics (Rx-type, non-medical use)	5.3	5.0	5.9	6.4	5.9	5.9	6.5	6.3	6.1	6.1	5.5
Pain Reliever	3.8	3.6	4.4	4.8	4.5	4.6	5.0	4.7	4.7	4.7	4.1
OxyContin®	0.4	0.5	0.7	0.5	0.4	0.5	0.4	0.4	0.4		
Stimulant	1.2	1.0	1.2	1.3	1.1	1.1	1.4	1.4	1.5	1.3	1.3
Methamphetamine	0.3	0.2	0.2	0.2	0.2	0.4	0.6	0.7	0.7	0.6	0.6
Sedative	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Tranquilizer	1.6	1.6	1.7	1.9	1.7	1.7	2.0	1.9	1.8	1.7	1.6

Estimated Lifetime Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	84.4	84.3	85.7	85.8	85.6	85.2	86.5	85.7	86.2	87.1	86.7
Any Tobacco	66.3	67.9	68.6	69.6	70.2	70.3	71.8	72.3	73.5	74.8	75.5
Cigarettes	59.5	61.0	62.1	63.7	64.2	64.7	66.6	67.3	68.7	70.2	71.2
Smokeless Tobacco	19.9	20.4	20.9	21.0	20.3	20.1	20.2	20.8	21.4	22.0	23.7
Any Illicit Drug	57.8	56.9	57.3	58.5	57.0	57.5	59.0	59.2	59.2	60.5	59.8
Marijuana	52.2	51.9	51.4	52.6	50.8	50.9	52.5	52.4	52.8	53.9	53.8
Inhalant	8.4	9.1	10.0	10.8	10.5	11.3	12.5	13.3	14.0	14.9	15.7
Hallucinogen	17.6	17.6	18.5	19.1	17.8	19.2	20.2	21.0	21.3	23.3	24.2
LSD	5.9	6.0	6.4	6.9	6.6	7.3	8.9	10.5	12.1	14.0	15.9
PCP	1.0	1.0	1.3	1.4	1.6	1.6	1.9	2.2	2.3	3.0	2.7
Ecstasy	12.9	12.3	12.4	12.5	12.2	12.8	13.4	13.7	13.8	14.8	15.1
Cocaine (any form)	12.3	12.4	13.4	14.9	14.5	15.0	15.7	15.1	15.2	15.0	15.4
Crack Cocaine	1.9	2.1	2.6	2.9	2.8	3.0	3.6	3.5	3.5	3.8	3.8
Heroin	1.9	1.7	1.8	1.7	1.4	1.5	1.6	1.5	1.6	1.6	1.6
Psychotherapeutics (Rx-type, non-medical use)	28.1	27.3	28.9	29.5	29.4	30.0	30.7	30.8	29.7	29.5	28.3
Pain Reliever	22.4	22.2	23.9	24.5	24.6	24.9	25.5	25.5	24.3	23.7	22.1
OxyContin®	5.8	6.1	6.2	6.2	5.3	5.2	5.1	5.0	4.3	3.6	2.6
Stimulant	9.8	9.2	10.4	10.7	10.6	10.9	12.0	12.3	12.1	12.2	12.2
Methamphetamine	3.1	3.4	4.0	4.5	4.7	5.6	6.4	6.7	7.0	6.9	7.4
Sedative	1.2	1.4	1.4	1.6	1.6	1.8	1.9	2.0	1.9	1.8	2.1
Tranquilizer	12.8	12.3	13.1	12.8	12.2	12.7	13.1	13.4	12.2	12.3	11.2
Needle/Injection Drug Use	1.4	1.2	1.1	1.3	0.9	1.3	1.2	1.4	1.3	1.3	1.2

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Some estimates from 2006-2010 differ slightly from those published in previous Status Reports due to recent SAMHSA updates.

Data Source:

U.S. Department of Health and Human Services, SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DetTabs/NSDUH-DetTabsTOC2012.htm>

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 26 AND OLDER, 2002-2012**

Estimated Past-Month Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	55.6	55.1	54.9	54.9	54.7	54.1	53.7	55.1	53.0	52.5	53.9
Binge Alcohol Use *	22.1	21.6	21.9	22.4	22.2	22.0	21.4	21.0	21.1	21.0	21.4
Heavy Alcohol Use **	6.1	5.7	6.1	6.2	6.3	6.1	6.0	5.6	6.1	5.9	5.9
Any Tobacco	27.0	26.3	27.2	27.3	28.4	28.6	29.4	29.0	28.5	29.3	29.9
Cigarettes	22.4	21.9	22.8	23.0	23.8	24.1	24.7	24.3	24.1	24.7	25.2
Smokeless Tobacco	3.3	3.0	3.1	3.1	3.3	3.0	3.1	3.0	2.7	3.2	3.2
Any Illicit Drug	7.0	6.3	6.6	6.3	5.9	5.8	6.1	5.8	5.5	5.6	5.8
Marijuana	5.3	4.8	4.8	4.6	4.2	3.9	4.2	4.1	4.1	4.0	4.0
Inhalant	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
Hallucinogen	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.2
LSD	0.0	0.0	***	0.0	***	0.0	0.0	0.0	0.0	0.0	0.0
PCP	0.0	***	0.0	0.0	***	0.0	0.0	0.0	0.0	***	0.0
Ecstasy	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Cocaine (any form)	0.6	0.4	0.5	0.6	0.7	0.7	0.8	0.8	0.7	0.8	0.7
Crack Cocaine	0.2	0.1	0.2	0.2	0.2	0.3	0.3	0.3	0.2	0.3	0.3
Heroin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1
Psychotherapeutics (Rx-type, non-medical use)	2.1	1.9	2.2	2.1	1.9	2.2	2.2	1.9	1.8	2.0	2.0
Pain Reliever	1.5	1.4	1.5	1.6	1.4	1.6	1.5	1.3	1.2	1.3	1.3
OxyContin®	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1		
Stimulant	0.3	0.3	0.3	0.4	0.2	0.3	0.4	0.3	0.4	0.4	0.4
Methamphetamine	0.2	0.2	0.1	0.2	0.1	0.2	0.3	0.2	0.2	0.3	0.2
Sedative	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2
Tranquilizer	0.7	0.6	0.7	0.7	0.6	0.6	0.5	0.6	0.5	0.6	0.6

Estimated Lifetime Substance Use (%)

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol	88.1	87.9	87.9	88.0	87.3	87.5	87.7	88.2	87.3	88.0	88.0
Any Tobacco	72.5	73.0	74.2	74.4	75.0	75.0	76.0	76.0	76.4	77.6	77.7
Cigarettes	67.9	68.6	70.0	70.1	70.7	70.9	71.7	71.9	72.3	73.6	73.7
Smokeless Tobacco	18.7	18.9	18.5	18.5	19.5	19.3	19.9	19.5	19.6	20.6	20.9
Any Illicit Drug	49.3	48.0	48.2	48.0	48.2	46.9	45.5	46.3	45.6	46.1	45.7
Marijuana	44.4	43.2	43.5	42.9	42.7	42.1	40.6	41.1	41.0	41.2	40.8
Inhalant	8.3	7.8	8.3	8.6	8.6	8.6	8.6	8.5	8.5	8.6	8.6
Hallucinogen	15.4	14.8	15.5	15.4	15.2	14.2	14.7	14.0	14.4	14.2	14.1
LSD	10.7	10.4	10.7	10.9	11.0	10.6	10.8	10.1	10.5	10.8	10.5
PCP	3.0	2.9	2.9	2.9	3.1	2.9	3.1	3.1	3.2	3.3	3.5
Ecstasy	5.6	4.9	5.7	4.9	4.4	4.1	3.9	3.6	3.4	3.1	2.6
Cocaine (any form)	16.5	16.3	16.6	16.2	16.5	16.1	15.7	15.2	15.6	16.3	15.9
Crack Cocaine	4.2	3.7	4.2	3.8	3.9	4.0	3.9	3.6	3.6	3.6	3.9
Heroin	1.9	1.8	1.8	1.6	1.7	1.7	1.7	1.6	1.4	1.7	1.7
Psychotherapeutics (Rx-type, non-medical use)	21.0	19.8	20.2	20.3	20.7	19.9	20.1	19.7	19.8	20.0	19.9
Pain Reliever	13.6	12.4	12.6	12.7	12.8	11.9	12.0	11.8	11.5	11.5	11.1
OxyContin®	2.1	1.7	1.9	1.7	1.4	1.2	1.1	0.9	0.8	0.8	0.5
Stimulant	8.8	8.4	9.1	9.2	8.9	9.1	9.4	8.7	9.6	10.0	10.4
Methamphetamine	5.5	5.3	5.9	5.8	5.7	5.8	6.3	5.5	6.5	6.9	7.1
Sedative	3.7	3.5	3.6	4.1	4.3	4.0	4.3	4.4	4.9	4.8	5.1
Tranquilizer	9.2	8.4	8.7	8.7	8.7	8.1	8.7	8.6	8.3	8.5	8.3
Needle/Injection Drug Use	1.8	1.6	1.6	1.7	2.0	1.9	1.8	1.7	1.6	1.8	2.0

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

*** Low precision, no estimate available

Some estimates from 2006-2010 differ slightly from those published in previous Status Reports due to recent SAMHSA updates.


Data Source:

U.S. Department of Health and Human Services, SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health.

<http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DetTabs/NSDUH-DetTabsTOC2012.htm>


ALCOHOL — PAST-MONTH USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12 and Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Alcohol Use (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL — PAST-MONTH BINGE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Binge Drinking (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL — PERCEIVED RISK OF BINGE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk of Binge Drinking Once or Twice Weekly (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL USE, BINGE USE, AND PERCEIVED RISK: MISSOURI AGE GROUP COMPARISONS

Estimated Past-Month Alcohol Use (%): Missouri Age Group Comparisons, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Binge Drinking (%): Missouri Age Group Comparisons, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perception of Great Risk of Binge Drinking (%): Missouri Age Group Comparisons, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL — PAST-MONTH, BINGE, AND PERCEIVED RISK OF USE: MISSOURI AND DBH REGIONS

Past-Month Alcohol Use (%): Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Past-Month Binge Drinking (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Perceptions of Great Risk of Binge Drinking (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


TOBACCO — PAST-MONTH CIGARETTE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Cigarette Use (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


TOBACCO — PERCEIVED RISK OF CIGARETTE USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12 & Older, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 12-17, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking 1+ Pack Cigarettes Daily (%): U.S. and Missouri Ages 18-25, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk Smoking One or More Packs of Cigarettes Daily (%): U.S. and Missouri Ages 26 & Older, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


TOBACCO USE AND PERCEIVED RISK: MISSOURI AGE GROUP COMPARISONS

Estimated Past-Month Tobacco Use (%): Missouri Age Group Comparisons, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Cigarette Use (%): Missouri Age Group Comparisons, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk Smoking One or More Packs of Cigarettes Daily (%): Missouri Age Group Comparisons, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


TOBACCO AND CIGARETTES — PAST-MONTH AND PERCEIVED RISK OF USE: MISSOURI AND DBH REGIONS

Estimated Past-Month Tobacco Use (%): Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Month Cigarette Use (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Perceive Great Risk in Smoking One or More Packs of Cigarettes Per Day (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


PAST-MONTH ILLICIT DRUG USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Use of Any Illicit Drug (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAST-MONTH ILLICIT DRUG USE EXCEPT MARIJUANA: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Month Use of Illicit Drug Other Than Marijuana (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Illicit Drug Other Than Marijuana (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Use of Illicit Drug Other Than Marijuana (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Use of Illicit Drug Other Than Marijuana (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ILLICIT DRUGS AND ILLICIT DRUGS EXCEPT MARIJUANA — PAST-MONTH USE: MISSOURI AND DBH REGIONS

Estimated Past-Month Illicit Drug Use (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Estimated Past-Month Use of Illicit Drugs Other Than Marijuana (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


MARIJUANA — PAST-MONTH USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Month Marijuana Use (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PAST-YEAR USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 12 & Older, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 12-17, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 18-25, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Marijuana Use (%): U.S. and Missouri Ages 26 & Older, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PERCEIVED RISK OF USE: U.S. AND MISSOURI COMPARISONS BY AGE GROUP

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Perceive Great Risk in Smoking Marijuana Once a Month (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAST MONTH, FIRST USE, AND PERCEIVED RISK OF MARIJUANA USE: MISSOURI AGE GROUPS

Estimated Past-Month Marijuana Use (%): Missouri Age Groups, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated First Use of Marijuana in Past Year (%): Missouri Age Groups, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Perception of Great Risk of Smoking Marijuana Once Per Month (%): Missouri Age Groups, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


MARIJUANA — PAST-MONTH, FIRST YEAR, AND PERCEIVED RISK OF USE: MISSOURI AND DBH REGIONS

Estimated Past-Month Marijuana Use (%): Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Average Annual Rate of First Use of Marijuana (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Estimated Perception of Great Risk of Smoking Marijuana Monthly (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


MARIJUANA, COCAINE, AND PAIN RELIEVERS — PAST-YEAR USE: MISSOURI AND DBH REGIONS

Estimated Past-Year Marijuana Use (%): Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Year Cocaine Use (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Estimated Past-Year Nonmedical Use of Pain Relievers (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


COCAINE — PAST-YEAR USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Cocaine Use (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAIN RELIEVERS: PAST-YEAR NON-MEDICAL USE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12 & Older, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 12-17, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 18-25, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Use of Pain Relievers (%): U.S. and Missouri Ages 26 & Older, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL AND ILLICIT DRUGS — PAST-YEAR DEPENDENCE OR ABUSE: U.S. AND MISSOURI AGE GROUPS

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%), U.S. and Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): U.S. and Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


PAST-YEAR ALCOHOL OR ILLICIT DRUG DEPENDENCE OR ABUSE: MISSOURI AGE GROUPS

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12 & Older, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 12-17, 2002-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 18-25, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Number with Past-Year Alcohol or Illicit Drug Dependence or Abuse: Missouri Ages 26 & Older, 2002-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


ALCOHOL AND ILLICIT DRUGS — PAST-YEAR SUBSTANCE DEPENDENCE OR ABUSE: MISSOURI AND DBH REGIONS

Estimated Past-Year Alcohol Dependence or Abuse (%): Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Year Illicit Drug Dependence or Abuse (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Estimated Past-Year Alcohol or Illicit Drug Dependence or Abuse (%): Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

PAST-YEAR SUBSTANCE DEPENDENCE OR ABUSE: MISSOURI DBH REGIONS BY AGE GROUP

Estimated Past-Year Substance Dependence or Abuse (%): Missouri and DBH Regions Ages 12-17, 2008-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Displayed percentages are derived from SAMHSA estimates for combined alcohol and illicit drug dependence/abuse.

Estimates are average annual rates based on combined years 2008-2010.

Estimated Past-Year Substance Dependence or Abuse (%): Missouri and DBH Regions Ages 18-25, 2008-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Displayed percentages are derived from SAMHSA estimates for combined alcohol and illicit drug dependence/abuse.

Estimates are average annual rates based on combined years 2008-2010.

Estimated Past-Year Substance Dependence or Abuse (%): Missouri and DBH Regions Ages 26 and Older, 2008-2010




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Displayed percentages are derived from SAMHSA estimates for combined alcohol and illicit drug dependence/abuse.

Estimates are average annual rates based on combined years 2008-2010.


PAST-YEAR UNMET NEED FOR SUBSTANCE ABUSE TREATMENT: MISSOURI AGE GROUPS

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 12 & Older, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 12-17, 2003-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 18-25, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


Estimated Number Needing but Not Receiving Treatment for Alcohol and Illicit Drugs: Missouri Ages 26 & Older, 2003-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.


UNMET NEED FOR SUBSTANCE ABUSE TREATMENT IN PAST YEAR: MISSOURI AND DBH REGIONS

Past-Year Percentage Needing but Not Receiving Treatment for Alcohol Use: Missouri and DBH Regions, 3-Year Spans



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Past-Year Percentage Needing but Not Receiving Treatment for Illicit Drug Use: Missouri and DBH Regions, 3-Year Spans



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

PAST-YEAR UNMET NEED FOR SUBSTANCE ABUSE TREATMENT: MISSOURI AND REGIONS BY AGE GROUP


Past-Year Unmet Need for Substance Abuse Treatment (%): Missouri and DBH Regions Ages 12-17, 2008-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are average annual rates based on combined years 2008-2010.


Past-Year Unmet Need for Substance Abuse Treatment (%): Missouri and DBH Regions Ages 18-25, 2008-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are average annual rates based on combined years 2008-2010.

Past-Year Unmet Need for Substance Abuse Treatment (%): Missouri and DBH Regions Ages 26 & Older, 2008-2010



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimates are average annual rates based on combined years 2008-2010.

ANY MENTAL ILLNESS IN PAST YEAR: U.S. AND MISSOURI COMPARISON BY AGE GROUP


Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 18 & Older, 2009-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 18-25, 2009-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

Estimated Past-Year Any Mental Illness: U.S. and Missouri Ages 26 and Older, 2009-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

SERIOUS MENTAL ILLNESS IN PAST YEAR: U.S. AND MISSOURI COMPARISON BY AGE GROUP


Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 18 & Older, 2009-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 18-25, 2009-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.

Estimated Past-Year Serious Mental Illness: U.S. and Missouri Ages 26 and Older, 2009-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Note: Due to adult mental health questionnaire changes in 2008, comparisons to prior years are not provided.


DEPRESSION — MAJOR PAST-YEAR EPISODE: U.S. AND MISSOURI COMPARISON BY AGE GROUP

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 12-17, 2005-2011



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18 and Older, 2005-2011



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.


Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 18-25, 2005-2011



Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.

Estimated Past-Year Major Depressive Episode (MDE): U.S. and Missouri Ages 26 and Older, 2005-2011




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health (NSDUH).

Note: Due to adult mental health questionnaire changes in 2008, MDE data beginning 2007/2008 are not comparable to prior years.


MAJOR DEPRESSIVE EPISODE IN PAST YEAR: MISSOURI AND DBH REGIONS

Estimated Past-Year Major Depressive Episode (MDE): Ages 12-17, Missouri and DBH Regions




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

Estimated Past-Year Major Depressive Episode (MDE): Ages 18-25, Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


Estimated Past-Year Major Depressive Episode (MDE): Ages 26 and Older, Missouri and DBH Regions



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.


SERIOUS THOUGHTS OF SUICIDE IN PAST YEAR: U.S. AND MISSOURI COMPARISON BY AGE GROUP AND MISSOURI ESTIMATES FOR DBH REGIONS

Estimated Past-Year Serious Thoughts of Suicide (%): U.S. and Missouri Ages 18 and Older, 2009-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Serious Thoughts of Suicide (%): U.S. and Missouri Ages 18-25, 2009-2011




Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Serious Thoughts of Suicide (%): U.S. and Missouri Ages 26 and Older, 2009-2011



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 2-year averages.

Estimated Past-Year Serious Thoughts of Suicide (%): Missouri and DBH Regions Ages 18 and Older, 2008-2010



Data Source: U.S. Dept of Health and Human Services, SAMHSA. National Survey on Drug Use and Health. Estimates are 3-year averages.

MISSOURI STUDENT SURVEY: PAST 30-DAY ALCOHOL, TOBACCO, AND OTHER DRUG USE

Reported Past 30-Day Substance Use (%): Missouri Students in Grades 6-12

		Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
Alcohol	2012	3.64	6.38	12.18	18.56	25.63	29.51	35.54
	2010	4.43	8.3	14.44	22.13	28.29	33.35	36.72
	2008	7.57	14.24	22.93	30.94	37.38	40.85	45.62
	2006	7.35	14.82	23.90	32.56	39.50	43.23	49.61
	2004	8.57	18.05	27.06	36.52	45.73	51.14	50.97
Alcohol: binge use in past 30 days	2012	1.04	2.61	6.05	10.20	16.13	19.69	24.18
	2010	1.60	3.50	7.10	12.37	17.34	22.45	24.59
Cigarettes	2012	2.57	4.36	8.23	11.38	15.13	20.06	21.84
	2010	4.00	5.92	10.94	14.80	18.68	22.95	24.99
	2008	3.02	6.65	9.68	14.03	18.97	22.56	24.34
	2006	3.77	7.28	11.82	14.85	19.87	21.63	28.04
	2004	5.27	10.45	15.22	18.30	22.72	30.80	28.30
Chewing Tobacco	2012	1.63	2.18	4.45	6.48	7.63	9.75	10.01
	2010	2.01	2.54	5.03	7.30	7.79	9.28	8.89
	2008	1.68	3.29	4.35	6.59	8.39	9.60	10.17
	2006	2.03	3.09	4.52	6.10	7.64	7.95	9.68
	2004	1.67	3.38	4.07	6.12	8.97	11.32	12.50
Marijuana	2012	1.00	2.32	5.97	9.55	13.05	15.51	21.35
	2010	0.99	2.29	5.50	10.25	15.64	18.31	19.52
	2008	0.83	3.11	5.54	8.97	12.90	14.86	16.56
	2006	0.83	2.72	5.05	9.32	13.57	15.53	17.50
	2004	0.95	3.34	7.23	10.53	13.06	13.03	14.60
Inhalant	2012	2.49	2.57	2.90	2.38	1.64	1.43	1.28
	2010	3.71	3.87	3.51	3.06	2.54	1.80	1.55
	2008	4.14	5.69	5.28	4.04	3.14	2.00	1.72
	2006	4.94	6.25	6.44	4.26	2.94	2.23	1.92
	2004	4.16	5.62	5.13	3.38	2.91	2.34	1.80
Pharmaceutical Drug: non-prescribed	2012	1.63	2.57	3.77	5.27	6.59	7.04	7.61
	2010	3.33	3.97	5.55	7.50	8.38	9.82	8.28
Over-the-Counter Drug: non-medical use	2012	1.31	2.18	3.50	4.08	5.08	4.68	3.81
	2010	2.14	3.07	4.39	5.97	6.19	6.05	5.27
Synthetic Drug	2012	1.49	1.67	2.41	3.31	3.95	4.26	3.52

Data Source: Missouri Institute of Mental Health. Missouri Student Survey.

Respondents participate voluntarily and do not comprise a scientifically selected sample.

2010 and 2012 data are weighted in accordance with population characteristics. 2004-2008 data are unweighted.

MISSOURI STUDENT SURVEY: LIFETIME ALCOHOL, TOBACCO, AND OTHER DRUG USE

Reported Lifetime Substance Use (%): Missouri Students in Grades 6-12

		Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
Alcohol	2012	13.25	19.96	31.97	42.76	52.80	59.30	64.53
	2010	15.09	24.39	36.49	47.99	55.42	63.20	65.33
	2008	30.07	44.52	56.04	65.36	70.76	73.60	74.71
	2006	35.24	48.41	61.47	69.79	75.37	77.12	82.39
Cigarettes	2012	7.23	10.78	19.02	25.02	31.07	37.94	41.20
	2010	8.35	12.54	22.11	28.62	36.42	43.43	45.46
	2008	9.53	17.57	24.06	30.80	38.02	43.34	45.48
	2006	20.40	27.38	34.60	39.39	46.70	49.71	55.50
Chewing Tobacco	2012	3.72	4.12	9.03	11.79	14.24	17.78	18.04
	2010	3.79	4.89	9.98	12.73	15.11	17.70	17.60
	2008	5.43	9.29	11.86	15.08	18.72	20.39	22.88
	2006	7.15	9.71	13.69	15.40	17.80	19.45	22.24
Marijuana	2012	1.73	4.49	10.56	18.17	25.64	31.74	40.66
	2010	1.85	4.42	10.44	18.76	28.73	35.35	38.86
	2008	2.02	6.49	11.83	19.01	27.28	32.69	35.72
	2006	2.41	6.35	11.69	19.83	30.17	34.78	40.69
Inhalants	2012	5.02	4.69	5.78	5.69	5.11	5.13	4.28
	2010	5.67	6.57	6.92	6.79	6.50	6.23	6.09
	2008	8.09	10.55	10.95	9.71	9.21	7.04	6.28
	2006	9.76	12.18	13.55	11.00	9.55	7.36	7.36
Heroin	2012	0.56	0.56	1.04	0.84	1.09	1.25	1.83
	2010	0.26	0.50	0.50	0.64	0.85	0.90	1.08
Cocaine *	2012	0.62	0.80	1.65	1.49	2.60	3.33	3.76
	2010	0.47	0.80	0.99	1.45	2.27	3.65	4.16
	2008	0.70	1.41	1.64	1.95	2.67	3.33	4.41
	2006	0.93	1.73	1.99	2.58	3.59	4.16	5.84
Methamphetamine **	2012	0.93	0.73	1.23	1.18	1.39	1.95	2.00
	2010	0.42	0.78	0.71	0.84	1.11	1.35	1.22
	2008	0.49	1.28	1.79	2.95	4.23	4.42	4.64
	2006	0.70	1.45	2.04	3.33	4.82	5.36	7.02
Hallucinogen	2012	0.62	1.14	2.18	2.66	4.10	5.29	7.52
	2010	0.46	0.97	1.85	3.19	4.43	6.36	5.90
	2008	0.39	0.85	1.40	2.22	3.59	4.53	5.23
	2006	0.40	0.94	1.34	2.11	3.27	3.89	5.31
Club Drug	2012	0.76	0.70	1.68	2.03	3.22	3.99	5.23
	2010	0.42	0.66	1.37	2.28	3.05	4.79	5.25
	2008	0.43	0.81	1.48	2.38	3.78	4.46	5.45
	2006	0.56	0.99	1.92	2.49	3.59	4.45	5.25
Steroid: non-prescribed	2012	1.07	1.08	1.83	1.39	1.34	1.62	1.71
	2010	0.92	0.86	0.99	1.24	1.28	1.14	0.85
	2008	1.62	2.29	2.55	2.60	2.68	2.59	2.64
	2006	1.44	2.05	2.52	2.55	2.60	2.74	2.92
Pharmaceutical Drug: non-prescribed	2012	3.06	4.20	6.77	9.28	11.93	14.16	16.12
	2010	5.23	6.56	9.08	12.13	13.96	16.93	15.96
Over-the-Counter Drug: non-medical use	2012	2.23	3.30	5.53	6.51	8.39	8.73	9.35
	2010	3.36	4.63	6.65	8.91	9.88	10.89	9.32
Synthetic Drug	2012	2.37	2.68	4.55	6.77	9.17	11.31	12.97

Data Source: Missouri Institute of Mental Health. Missouri Student Survey.

Respondents participate voluntarily and do not comprise a scientifically selected sample.


2010 and 2012 data are weighted in accordance with population characteristics. 2006-2008 data are unweighted.

* Item specified "crack or freebase form of cocaine" in 2006-2008 surveys and "any form of cocaine" in 2010-2012 surveys.

** Item specified "speed, amphetamine, or methamphetamine" in 2002-2008 surveys and "methamphetamine" in 2010-2012 surveys.


ALCOHOL USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Alcohol (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Alcohol (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 2-Week (2004-2008) and Past 30-Day (2010-2012) Binge Alcohol Use (%): Missouri Students in Grades 6-12



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


TOBACCO USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Cigarettes (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Cigarettes (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Lifetime Use of Chewing Tobacco (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Chewing Tobacco (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


MARIJUANA USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Marijuana (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Marijuana (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


COCAINE AND HEROIN USE: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Cocaine (%): Missouri Students in Grades 6-12, 2004-2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.

Past 30-Day Use of Cocaine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


Lifetime Use of Heroin (%): Missouri Students in Grades 6-12, 2010-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


METHAMPHETAMINE USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Speed, Amphetamine, or Methamphetamine (%): Missouri Students in Grades 6-12, 2004-2008




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.

Lifetime Use of Methamphetamine (%): Missouri Students in Grades 6-12, 2010-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Past 30-Day Use of Speed, Amphetamine or Methamphetamine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


INHALANT USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Inhalants (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Inhalants (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


HALLUCINOGEN USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Hallucinogens such as LSD, PCP, Mescaline, Peyote (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of LSD or other Psychedelics (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


CLUB DRUG USE IN LIFETIME AND PAST 30-DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Club Drugs such as Ecstasy, GHB, Rohypnol, Ketamine (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


Past 30-Day Use of Club Drugs such as Ecstasy, GHB, Rohypnol, Ketamine (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are unweighted.


PHARMACEUTICAL DRUG USE IN LIFETIME AND PAST 30 DAYS: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Non-Prescribed Pharmaceutical Drugs (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Past 30-Day Use of Non-Prescribed Pharmaceutical Drugs (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Lifetime Use of Non-Prescribed Steroids (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


OVER-THE-COUNTER AND SYNTHETIC DRUG USE: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Use of Over-the-Counter Medicines for Non-medical Reason (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Past 30-Day Use of Over-the-Counter Meds for Non-medical Reason (%): Missouri Students in Grades 6-12, 2012




Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

Lifetime Use of Synthetic Drugs such as K2, Bath Salts, Spice (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.


Past 30-Day Use of Synthetic Drugs such as K2, Bath Salts, Spice (%): Missouri Students in Grades 6-12, 2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. Data are weighted.

LIFETIME INJECTION DRUG USE: MISSOURI STUDENTS IN GRADES 6-12

Lifetime Injection Drug Use (%): Missouri Students in Grades 6-12, 2004-2008



Data Source: Missouri Institute of Mental Health. 2004-2008 Missouri Student Survey respondents. Data are unweighted.

**YOUTH RISK BEHAVIOR SURVEILLANCE SYSTEM (YRBSS)
TABLES FOR ALCOHOL, TOBACCO, AND PRESCRIPTION DRUG USE**

	2011		2009		2007		2005		2003		2001		1999		1997	
	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.
Alcohol Use in Lifetime (%)																
Overall Grades 9-12	---	70.8	70.5	72.5	73.5	75.0	72.1	74.3	77.8	74.9	80.1	78.2	79.4	81.0	78.3	79.1
Grade 9	---	61.7	62.0	63.4	67.8	65.5	64.5	66.5	73.6	65.0	76.4	73.1	73.6	73.4	69.3	72.0
Grade 10	---	69.2	68.3	71.1	74.1	74.7	70.6	74.4	75.9	75.7	78.8	76.3	79.2	83.2	82.1	77.4
Grade 11	---	75.3	76.8	77.8	77.7	79.4	75.7	76.3	81.0	78.6	78.6	80.4	84.6	80.8	77.0	81.9
Grade 12	---	79.0	77.5	79.7	75.9	82.8	79.2	81.7	81.9	83.0	87.7	85.1	80.8	88.3	86.1	84.0
Alcohol Use in Past 30 Days (%)																
Overall Grades 9-12	---	38.7	39.3	41.8	44.4	44.7	40.8	43.3	49.2	44.9	47.6	47.1	49.9	50.0	51.4	50.8
Grade 9	---	29.8	28.6	31.5	38.5	35.7	30.4	36.2	44.4	36.2	43.2	41.1	45.4	40.6	42.6	44.2
Grade 10	---	35.7	37.7	40.6	44.7	41.8	41.5	42.0	46.1	43.5	47.7	45.2	43.3	49.7	50.9	47.2
Grade 11	---	42.7	48.4	45.7	46.7	49.0	44.3	46.0	50.9	47.0	48.4	49.3	55.5	50.9	52.3	53.2
Grade 12	---	48.4	45.8	51.7	48.6	54.9	48.8	50.8	57.3	55.9	52.2	55.2	57.2	61.7	62.2	57.3
Binge Drinking in Past 30 Days (%)																
Overall Grades 9-12	---	21.9	25.3	24.2	29.1	26.0	24.9	25.5	30.5	28.3	34.1	29.9	32.0	31.5	37.9	33.4
Grade 9	---	14.0	15.2	15.3	22.3	17.0	13.6	19.0	25.3	19.8	28.0	24.5	23.5	21.1	28.4	25.7
Grade 10	---	18.4	24.0	22.3	26.8	23.7	26.3	24.6	27.6	27.4	33.0	28.2	28.2	32.2	35.5	29.9
Grade 11	---	25.2	32.1	28.3	33.0	29.9	29.0	27.6	32.0	31.8	36.5	32.2	40.8	34.0	39.1	37.5
Grade 12	---	31.5	32.7	33.5	35.3	36.5	32.4	32.8	38.9	37.2	40.0	36.7	38.7	41.6	51.3	39.3
Alcohol Use Before Age 13 (%)																
Overall Grades 9-12	---	20.5	20.4	21.1	20.9	23.8	24.2	25.6	25.4	27.8	30.0	29.1	33.5	32.2	33.6	31.1
Grade 9	---	26.6	23.9	28.1	30.5	30.9	34.1	33.9	38.2	36.4	38.7	39.7	40.2	40.4	42.4	41.9
Grade 10	---	21.1	22.9	22.2	20.5	24.4	22.5	26.2	26.4	28.5	31.2	28.8	38.0	35.6	32.6	32.0
Grade 11	---	17.6	18.2	17.9	17.8	19.6	23.8	20.5	20.3	23.0	26.5	23.4	30.6	26.2	29.1	29.9
Grade 12	---	15.1	15.3	14.2	11.8	18.0	14.4	19.3	14.0	20.3	21.3	21.2	21.4	24.3	27.3	22.8
Cigarette Use in Lifetime (%)																
Overall Grades 9-12	---	44.7	46.5	46.3	50.9	50.3	49.6	54.3	63.3	58.4	68.5	63.9	69.8	70.4	74.0	70.2
Grade 9	---	37.6	34.0	37.7	47.2	42.7	40.7	48.7	60.8	52.0	62.5	58.4	67.7	61.8	72.3	67.7
Grade 10	---	41.0	44.1	44.0	45.9	48.8	51.2	52.5	59.7	58.3	68.9	62.6	70.8	73.9	73.7	70.0
Grade 11	---	47.1	52.9	50.0	57.4	53.4	53.4	57.5	66.6	60.0	68.4	65.9	69.5	69.9	73.0	68.8
Grade 12	---	54.5	57.9	55.5	53.7	59.3	54.3	60.3	66.9	65.4	75.9	71.1	72.1	78.0	77.4	73.7
Cigarette Use in Past 30 Days (%)																
Overall Grades 9-12, U.S.	---	18.1	18.9	19.5	23.8	20.0	21.3	23.0	24.8	21.9	30.3	28.5	32.8	34.8	40.3	36.4
Grade 9, U.S.	---	13.0	9.4	13.5	22.2	14.3	14.1	19.7	25.6	17.4	27.0	23.9	27.1	27.6	34.9	33.4
Grade 10, U.S.	---	15.6	19.0	18.3	17.8	19.6	21.4	21.4	23.5	21.8	29.8	26.9	33.9	34.7	39.1	35.3
Grade 11, U.S.	---	19.3	22.8	22.3	30.6	21.6	24.3	24.3	28.2	23.6	29.6	29.8	38.1	36.0	40.1	36.6
Grade 12, U.S.	---	25.1	26.7	25.2	25.4	26.5	26.5	27.6	21.4	26.2	35.4	35.2	33.5	42.8	48.2	39.6
Cigarette Use Before Age 13 (%)																
Overall Grades 9-12, U.S.	---	10.3	9.9	10.7	13.4	14.2	14.8	16.0	20.2	18.3	25.7	22.1	25.2	24.7	27.4	24.8
Grade 9, U.S.	---	12.1	8.2	12.1	15.7	16.3	15.9	18.6	28.1	19.3	31.1	26.2	30.4	27.0	34.2	32.0
Grade 10, U.S.	---	10.1	13.3	11.2	12.6	14.3	14.2	16.0	21.9	20.1	26.5	22.9	27.2	28.5	24.3	27.5
Grade 11, U.S.	---	9.8	8.3	10.3	14.3	12.0	16.6	14.4	17.4	16.0	22.2	18.5	23.2	21.1	25.4	22.2
Grade 12, U.S.	---	8.2	9.7	8.6	10.1	13.3	12.3	13.9	11.3	16.5	21.1	19.0	17.8	20.7	23.3	18.6
Prescription Drug Use without Rx in Lifetime (%)																
Overall Grades 9-12, U.S.	---	20.7														
Grade 9, U.S.	---	16.5														
Grade 10, U.S.	---	18.2														
Grade 11, U.S.	---	23.3														
Grade 12, U.S.	---	25.6														


Values shown are weighted estimates based on sampling. Apparent differences in values may not be statistically significant.

--- 2011 Missouri estimates are not available due to insufficient response rate to weight data.

Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention.


PAST 30-DAY ALCOHOL USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Alcohol Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey.

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.


PAST 30-DAY BINGE ALCOHOL USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey.

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.


PAST 30-DAY, BINGE, AND EARLY ALCOHOL USE: MISSOURI GRADE 9-12 COMPARISONS, 1995-2009

Past 30-Day Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Binge Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Alcohol Use Before Age 13 (%): Missouri Students in Grades 9-12, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.


PAST 30-DAY CIGARETTE USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Dept. of HHS, Centers for Disease Control and Prevention. Estimates from the Youth Risk Behavior Survey

Past 30-Day Cigarette Use (%): U.S. and Missouri Students in Grade 12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey.

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.

**YOUTH RISK BEHAVIOR SURVEILLANCE SYSTEM (YRBSS)
TABLES FOR ILLICIT DRUG USE**

	2011		2009		2007		2005		2003		2001		1999		1997	
	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.	MO	U.S.
Marijuana Use in Past 30 Days (%)																
Overall Grades 9-12	NA	23.1	20.6	20.8	19.0	19.7	18.1	20.2	21.8	22.4	24.4	23.9	25.6	26.7	28.2	26.2
Grade 9	NA	18.0	13.7	15.5	12.5	14.7	11.7	17.4	20.7	18.5	25.8	19.4	22.3	21.7	23.4	23.6
Grade 10	NA	21.6	17.6	21.1	16.7	19.3	22.2	20.2	18.0	22.0	21.9	24.8	24.5	27.8	26.2	25.0
Grade 11	NA	25.5	27.6	23.2	26.6	21.4	23.2	21.0	23.0	24.1	24.8	25.8	31.8	26.7	29.9	29.3
Grade 12	NA	28.0	25.0	24.6	21.2	25.1	15.5	22.8	25.4	25.8	24.6	26.9	24.5	31.5	34.9	26.6
Marijuana Use Before Age 13 (%)																
Overall Grades 9-12	NA	8.1	6.7	7.5	7.8	8.3	8.8	8.7	10.0	9.9	12.4	10.2	13.6	11.3	9.8	9.7
Grade 9	NA	9.7	6.3	9.1	8.9	9.8	11.4	11.2	13.1	11.7	15.2	11.6	17.0	12.7	14.3	14.9
Grade 10	NA	7.5	6.7	8.3	7.1	8.7	8.5	9.1	11.7	10.8	10.9	12.1	13.1	12.6	7.9	10.4
Grade 11	NA	7.6	7.1	6.5	8.9	7.2	9.5	7.1	7.1	8.1	12.9	8.5	14.8	9.5	7.8	8.3
Grade 12	NA	7.0	6.7	5.2	5.5	6.6	5.2	6.2	6.4	7.8	9.5	7.8	8.2	9.5	7.6	5.8
Ecstasy Use in Lifetime (%)																
Overall Grades 9-12	NA	8.2	5.7	6.7	6.9	5.8	6.1	6.3	6.7	11.1	NA	11.1				
Grade 9	NA	5.2	1.3	4.9	5.5	4.6	3.8	5.8	5.9	10.9	NA	7.5				
Grade 10	NA	7.7	6.2	5.2	5.1	5.3	5.2	6.0	4.5	9.0	NA	10.5				
Grade 11	NA	9.2	7.5	8.7	8.2	5.6	7.3	6.5	7.2	11.4	NA	13.7				
Grade 12	NA	11.3	8.4	8.0	8.2	7.6	8.5	6.7	9.7	12.8	NA	14.1				
Inhalant Use in Lifetime (%)																
Overall Grades 9-12	NA	11.4	10.2	11.7	12.7	13.3	12.2	12.4	10.5	12.1	12.7	14.7	12.8	14.6	17.4	16.0
Grade 9	NA	12.7	10.7	13.0	17.3	15.0	12.5	14.1	13.5	13.6	13.9	17.4	13.8	16.5	18.4	19.0
Grade 10	NA	11.8	11.1	12.5	11.0	14.6	13.1	13.2	11.2	11.1	14.2	14.0	13.8	16.0	19.9	16.5
Grade 11	NA	11.1	9.3	11.5	10.4	12.5	10.8	11.4	10.5	11.0	12.1	13.8	12.3	13.4	13.2	15.4
Grade 12	NA	9.3	9.1	9.1	10.1	10.2	11.3	10.1	6.2	11.8	9.6	12.5	9.8	11.3	16.6	13.4
Cocaine Use in Lifetime (%)																
Overall Grades 9-12	NA	6.8	5.0	6.4	6.6	7.2	7.4	7.6	6.6	8.7	8.6	9.4	7.7	9.5	9.3	8.2
Grade 9	NA	5.0	2.2	4.5	4.5	4.8	3.1	6.0	5.4	6.8	8.1	7.2	6.7	5.8	6.0	6.7
Grade 10	NA	6.5	4.7	5.6	4.6	7.2	7.2	7.2	7.1	8.5	5.0	8.6	6.0	9.9	7.3	7.5
Grade 11	NA	7.5	6.8	7.7	8.2	7.7	8.8	8.7	7.1	9.0	10.2	10.4	8.6	9.9	9.6	9.1
Grade 12	NA	8.5	6.8	7.9	9.4	9.5	11.1	8.9	7.0	10.5	11.4	12.1	9.3	13.7	14.5	9.2
Methamphetamine Use in Lifetime (%)																
Overall Grades 9-12	NA	3.8	3.7	4.1	3.7	4.4	6.4	6.2	6.2	7.6	10.4	9.8	8.2	9.1		
Grade 9	NA	3.2	1.9	3.3	2.9	3.6	2.9	5.7	5.1	6.7	9.7	8.1	5.7	6.3		
Grade 10	NA	3.7	3.3	3.7	2.3	4.1	4.9	5.9	8.0	7.5	7.5	9.7	8.2	9.3		
Grade 11	NA	4.1	4.9	5.2	4.6	5.4	8.4	6.7	6.6	8.0	9.7	9.2	8.4	10.1		
Grade 12	NA	4.1	4.3	4.1	4.7	4.5	9.7	6.4	5.0	8.0	14.8	12.8	10.8	11.5		
Heroin Use in Lifetime (%)																
Overall Grades 9-12	NA	2.9	2.8	2.5	2.3	2.3	2.7	2.4	1.7	3.3	2.7	3.1	2.1	2.4		
Grade 9	NA	2.9	0.9	2.1	2.0	2.6	2.1	2.8	3.4	3.5	2.5	3.2	1.8	2.0		
Grade 10	NA	2.8	2.5	2.2	1.8	1.8	2.0	2.5	1.2	2.9	1.5	3.3	2.4	1.8		
Grade 11	NA	2.8	3.6	3.2	2.5	1.8	3.0	1.8	0.9	3.0	2.5	2.8	1.5	2.6		
Grade 12	NA	2.7	4.2	2.5	2.1	2.6	3.2	2.0	1.0	2.9	3.7	3.0	1.9	3.3		
Injection Drug Use in Lifetime (%)																
Overall Grades 9-12	NA	2.3	2.4	2.1	2.5	2.0	2.3	2.1	1.3	3.2	1.6	2.3	2.0	1.8	2.8	2.1
Grade 9	NA	2.1	1.3	2.0	3.6	2.0	1.7	2.4	3.2	3.2	1.5	2.5	2.4	1.6	3.5	3.0
Grade 10	NA	2.3	2.6	2.0	1.8	1.4	1.4	2.3	0.2	3.2	1.2	2.6	1.6	1.2	1.5	2.5
Grade 11	NA	2.4	2.0	2.5	2.3	1.9	2.9	1.7	0.9	2.8	1.4	1.9	1.6	2.0	2.5	1.6
Grade 12	NA	2.2	3.6	1.8	1.7	2.4	2.7	1.7	0.5	3.0	1.8	2.1	1.7	2.3	3.0	1.5


Values shown are weighted estimates based on sampling. Apparent differences in values may not be statistically significant.

--- 2011 Missouri estimates are not available due to insufficient response rate to weight data.

Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention.


PAST 30-DAY MARIJUANA USE: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Estimated Past 30-Day Marijuana Use (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated Past 30-Day Marijuana Use (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated Past 30-Day Marijuana Use (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated Past 30-Day Marijuana Use (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey.

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.


FIRST USE OF MARIJUANA BEFORE AGE 13: U.S. AND MISSOURI STUDENTS IN GRADES 9-12, 1995-2009

Estimated First Use of Marijuana Before Age 13 (%): U.S. and Missouri Students in Grade 9, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated First Use of Marijuana Before Age 13 (%): U.S. and Missouri Students in Grade 10, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated First Use of Marijuana Before Age 13 (%): U.S. and Missouri Students in Grade 11, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Estimated First Use of Marijuana Before Age 13 (%): U.S. and Missouri Students in Grade 12, 1995-2009




Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey.

Data shown are weighted estimates. 2011 Missouri estimates are not available due to insufficient response rate to weight data.


PAST YEAR, PAST 30-DAY, AND BINGE ALCOHOL USE AMONG MISSOURI COLLEGE STUDENTS

Past-Year Alcohol Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents.

Past 30-Day Alcohol Use (%): Students Completing the Missouri College Health Behavior Survey




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents.

Past 2-Week Binge Alcohol Use in One Sitting (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents.


Past 2-Week Binge Alcohol Use in 2-Hour Period (%): Students Completing the Missouri College Health Behavior Survey



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents.


PAST-YEAR ILLICIT DRUG USE AMONG MISSOURI COLLEGE STUDENTS

Past-Year Marijuana Use (%): Students Completing the Missouri College Health Behavior Survey, 2007-2013




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2012 respondents.

Past-Year Cocaine Use (%): Students Completing the Missouri College Health Behavior Survey, 2007-2013




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents. Item not included on 2012 survey.

Past-Year Amphetamine Use (%): Students Completing the Missouri College Health Behavior Survey, 2007-2013




Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents. Item not included on 2012 survey.

Past-Year Inhalant Use (%): Students Completing the Missouri College Health Behavior Survey, 2007-2013



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents. Item not included on 2012 survey.

Past-Year Club Drug Use (%): Students Completing the Missouri College Health Behavior Survey, 2007-2013



Data Source: University of Missouri--Columbia. Partners in Prevention Statewide Coalition.
Missouri College Health Behavior Survey 2007-2013 respondents. Item not included on 2012 survey.

MISSOURI COLLEGE HEALTH BEHAVIOR SURVEY

Reported Past-Year Substance Involvement among Missouri College Students

	2013	2012	2011	2010	2009	2008
ALCOHOL AND DRUG USE (%)						
Alcohol Use in Past Year	77.0	79.3	77.8	79.9	79.5	81.3
Alcohol Use in Past 30 Days	63.5	62.6	66.2	67.2	66.7	68.1
Binge Alcohol Use in One Sitting in Past 2 Weeks	27.2	27.9	32.8	33.8	35.7	31.2
Binge Alcohol Use in 2-Hour Period in Past 2 Weeks	25.6	26.3	31.2	31.7	33.5	29.3
Marijuana Use in Past Year	12.6	21.0	23.4	21.6	23.1	21.8
Cocaine Use in Past Year	2.2	---	2.4	2.3	2.9	3.4
Amphetamine Use in Past Year	1.9	---	3.3	2.4	2.2	2.1
Methamphetamine Use in Past Year	1.0	---	0.8	0.5	0.7	0.8
Inhalant Use in Past Year	1.1	---	1.2	0.9	0.7	0.8
Club Drug Use in Past Year	2.9	---	2.8	2.5	2.6	2.4
K2 (Synthetic Marijuana) Use in Past Year	1.9	3.2	---	---	---	---
Bath Salts Use in Past Year	1.1	1.0	---	---	---	---
Prescription Stimulant Use without Rx in Past Year	6.6	5.4	5.5	4.6	5.3	4.6
Prescription Pain Medication Use without Rx in Past Year	7.6	4.5	6.6	6.5	8.4	8.2
Prescription Sedative Use without Rx in Past Year	2.7	2.1	2.8	2.6	3.8	3.4
Prescription Sleep Medication Use without Rx in Past Year	2.0	1.0	1.8	2.0	2.5	2.2

AMONG ALL STUDENTS, DRINKING RELATED RISKS AND CONSEQUENCES EXPERIENCED (%)

Hangover	50.8	48.6	51.4	52.4	53.0	55.1
Rode with Someone who Drove After Drinking	30.6	29.8	33.6	36.3	36.9	38.7
Blackout or Memory Loss	35.6	26.2	26.1	25.1	25.7	27.7
Driven After Consumption	25.2	25.5	27.3	30.0	30.5	31.9
Missed Class	16.9	16.5	18.3	20.4	21.5	24.3
Engaged in Risky Sexual Behavior	16.2	16.5	16.9	17.2	18.3	19.5
Forced or Coerced to Drink More than Desired	17.3	16.1	13.5	12.6	13.9	---
Hurt or Injured	11.5	11.4	11.6	11.6	11.6	12.5
Performed Poorly on Test or Assignment	10.4	10.0	11.0	12.2	12.5	14.2
Taken Advantage of Sexually	7.1	6.3	6.3	5.7	6.3	6.2
Trouble with Campus Administration	3.7	3.0	3.5	2.0	2.9	3.1
Took Advantage of Someone Sexually	2.6	2.4	2.2	2.0	2.3	---
Received Medical Attention	2.6	2.2	2.1	2.2	2.5	2.3
Arrested by Campus Police or Other Law Enforcement	2.3	1.9	1.8	2.3	2.3	2.0
Arrested for DUI or DWI	1.6	0.8	1.1	1.4	1.2	1.0

AMONG PAST-YEAR DRINKERS, THOSE RESPONDING "ALWAYS" OR "USUALLY" TO USE OF THE FOLLOWING PROTECTIVE BEHAVIORS (%)

Know Where One's Drink Has Been At All Times	80.9	81.2	82.1	83.4	---	---
Eat Before and/or During Alcohol Consumption	76.6	77.5	75.5	76.4	77.2	76.6
Go Home with a Friend	65.3	65.0	65.1	65.5	---	---
Use a Designated Driver	64.3	64.8	66.3	69.9	70.7	70.7
Purposefully Limit the Amount of Money Spent on Alcohol	61.4	61.2	60.3	63.8	61.0	59.5
Avoid Trying to "Keep-up" or "Out-drink" Others	60.1	58.4	59.7	59.1	---	---
Drink Slowly, Rather than Gulp or Chug	53.6	48.4	49.1	49.2	---	---
Determine Not to Exceed a Set Number of Drinks	40.6	40.9	43.0	45.0	44.4	43.7
Avoid Mixing Different Types of Alcohol	41.7	38.3	38.4	39.4	---	---
Drink Water While Drinking Alcohol	36.6	35.9	34.4	33.2	---	---
Avoid Drinking Games	31.5	33.5	32.5	37.2	34.8	36.4
Alternate Alcoholic and Non-alcoholic Drinks	31.4	30.4	30.4	30.1	29.2	28.0
Have a Friend Tell You When You Have Had Enough	28.3	29.1	29.0	32.6	34.8	36.4
Leave the Bar or Party at Predetermined Time	28.6	29.1	28.2	30.0	---	---
Stop Drinking at a Predetermined Time	27.7	26.4	26.1	26.1	---	---
Put Extra Ice in Drink	21.8	20.4	20.0	20.3	---	---

Data Source:

Partners in Prevention Statewide Coalition.

<http://pip.missouri.edu/>

Based on responses of 9599 students at 21 Missouri colleges in 2013, 9151 students at 21 colleges in 2012, 8905 students at 20 colleges in 2011, 6543 students at 13 colleges in 2010, 6257 students at 13 colleges in 2009, and 5050 students at 12 colleges in 2008.

MISSOURI COLLEGE HEALTH BEHAVIOR SURVEY

Reported Past-Year Mental Health Problems among Missouri College Students

	2013	2012	2011	2010	2009	2008
LEVEL OF STRESS EXPERIENCED IN PAST TWO WEEKS (%)						
None	1.9	1.9	1.7	2.0	2.1	---
Minimal	10.7	11.1	10.5	13.9	11.6	---
A Little	18.7	19.0	18.8	20.4	20.2	---
Manageable	47.5	47.7	47.2	44.6	45.4	---
Overwhelming	18.3	18.1	19.5	16.7	18.5	---
Unbearable	2.1	2.0	2.3	2.1	2.3	---
AMONG ALL STUDENTS, MAIN SOURCES OF STRESS (%)						
School / Academics	---	77.2	78.7	75.2	74.4	86.8
Financial Concerns	---	47.4	50.6	49.2	50.3	50.8
Future Plans	---	43.9	44.3	42.9	50.2	---
Time Management	---	43.1	47.1	43.6	52.8	49.0
Job	---	31.1	32.1	32.9	34.1	34.9
Dating / Relationship with Partner	---	26.2	28.8	31.8	36.0	34.0
Family	---	25.0	27.7	28.4	31.6	25.1
Physical Health	---	20.1	22.3	23.3	22.2	21.0
Outside Organizations / Responsibilities	---	17.7	17.7	15.5	18.1	---
Friends	---	16.1	17.6	18.5	22.3	17.1
Roommates	---	14.6	15.8	15.3	17.7	17.9
Mental Health Issues	---	7.2	7.4	8.1	8.9	---
Other Source of Stress	---	2.6	3.2	3.7	4.1	5.6
AMONG ALL STUDENTS, MENTAL HEALTH PROBLEMS EXPERIENCED IN PAST YEAR (%)						
Anxiety	39.4	34.4	34.9	32.7	35.8	35.2
Chronic Sleep Disorder	18.1	15.0	16.8	15.6	16.6	19.4
Suicidal Thoughts	13.6	13.6	13.4	13.1	---	---
Major Depression	17.6	12.5	12.5	12.1	12.9	35.2
Panic Attack	13.4	11.7	12.5	11.6	12.3	---
Eating Disorder	6.4	5.3	5.0	4.7	6.0	6.3
Self-injury	3.8	3.5	4.0	3.7	3.9	2.4
Abusive Relationship	3.1	3.0	3.2	2.8	3.1	---
Alcohol Abuse/dependency	3.2	2.8	2.8	3.1	4.3	3.0
Bipolar Disorder	2.3	2.1	2.3	2.3	2.7	---
Sexual Assault	1.4	1.1	1.2	0.9	1.1	1.2
Suicide Attempt	0.9	1.0	0.8	0.9	---	---
Other Problem	1.7	1.5	1.5	1.8	1.8	---
None of the Listed Problems	46.7	42.2	42.3	43.7	45.7	38.8
AMONG STUDENTS EXPERIENCING PROBLEMS, THOSE WHO SOUGHT ASSISTANCE FOR THE FOLLOWING (%)						
Major Depression	48.4	53.3	50.0	48.5	47.5	59.3
Panic Attack	44.4	43.1	41.1	38.3	39.6	---
Bipolar Disorder	49.2	42.0	51.5	48.7	44.7	---
Abusive Relationship	33.9	38.5	35.8	34.8	32.0	---
Anxiety	37.1	36.8	34.2	32.2	33.2	30.1
Chronic Sleep Disorder	31.2	32.6	31.1	29.7	29.8	24.7
Self-injury	38.1	30.9	32.8	28.5	31.0	40.7
Sexual Assault	27.9	28.2	26.6	32.2	22.1	35.5
Eating Disorder	26.7	27.2	23.1	30.2	23.1	27.2
Alcohol Abuse/dependency	18.4	22.6	16.6	21.2	19.0	28.0
Other Problem	45.6	48.6	45.4	46.2	45.5	---

Data Source:

Partners in Prevention Statewide Coalition.

<http://pip.missouri.edu/>

Based on responses of 9599 students at 21 Missouri colleges in 2013, 9151 students at 21 colleges in 2012, 8905 students at 20 colleges in 2011, 6543 students at 13 colleges in 2010, 6257 students at 13 colleges in 2009, and 5050 students at 12 colleges in 2008.

MISSOURI COLLEGE HEALTH BEHAVIOR SURVEY

Reported Gambling among Missouri College Students

	2013	2012	2011	2010	2009	2008
INVOLVEMENT IN GAMBLING (%)						
Yes	27.9	28.7	29.9	31.1	21.6	28.2
No	70.9	71.3	70.1	68.9	78.4	71.8

AMONG STUDENTS WHO GAMBLE, THOSE RESPONDING "SOMETIMES", "OFTEN", OR "ALWAYS" TO THE FOLLOWING AS A RESULT OF GAMBLING (%)

Lost Track of Time While Gambling	---	11.8	---	12.5	16.9	7.4
Spent Time Thinking About Past Gambling Experiences	---	10.2	---	10.2	14.1	---
Spent Time Planning Future Gambling Activities	---	8.9	---	8.5	14.5	---
Lost a Lot of Time	---	7.1	---	6.9	8.0	3.7
Attempted to Recoup Gambling Losses	---	6.9	---	7.6	9.6	---
Felt Need to Gamble Increased Amounts for Excitement	---	6.9	---	6.5	8.5	---
Sought Ways to Find Money to Gamble	---	5.7	---	2.5	7.4	---
Lied to Family or Friends to Hide Gambling Activity	---	5.2	---	5.0	6.3	---
Withdrew from Social Situations or Enjoyable Activities	---	5.1	---	4.8	7.1	---
Strained Relationships	---	5.0	---	4.6	5.8	---
Borrowed Money for Gambling Debts	---	4.9	---	4.5	5.9	---

AMONG STUDENTS WHO GAMBLE, THOSE RESPONDING "SOMEWHAT", "QUITE A BIT", OR "TO A GREAT EXTENT" TO THE FOLLOWING DUE TO GAMBLING (%)

Lost a Considerable Amount of Money	---	6.6	---	6.6	8.0	4.1
Gambled Money Intended for Something Else	---	4.8	---	5.2	7.4	3.8
Experienced Financial Hardship	---	4.6	---	5.4	6.4	2.9
Academic Problems	---	2.7	---	3.0	3.0	2.2
Legal Trouble	---	2.0	---	1.7	2.1	0.9

Data Source:

University of Missouri--Columbia. Partners in Prevention Statewide Coalition.

<http://pip.missouri.edu/>

Based on responses of 9599 students at 21 Missouri colleges in 2013, 9151 students at 21 colleges in 2012, 8905 students at 20 colleges in 2011, 6543 students at 13 colleges in 2010, 6257 students at 13 colleges in 2009, and 5050 students at 12 colleges in 2008.

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (BRFSS)
TABLES FOR ALCOHOL, TOBACCO, AND MENTAL HEALTH: AGES 18 AND OLDER**

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002
Alcohol Use in Past 30 Days (%)											
Missouri	48.5	52.5	47.7	50.1	51.9	51.9	50.7	51.8	51.8	53.5	53.6
U.S. (Median of the States)	55.3	57.1	54.6	54.3	54.4	54.8	55.4	56.2	57.1	59.4	58.1
Binge Drinking * in Past 30 Days (%)											
Missouri	17.3	19.2	15.6	17.2	15.3	16.2	16.5				
U.S. (Median of the States)	16.9	18.3	15.1	15.7	15.6	15.8	15.4				
* 5 or more drinks on one occasion for males											
* 4 or more drinks on one occasion for females											
Heavy Drinking ** in Past 30 Days (%)											
Missouri	6.3	7.3	5.5	4.1	5.3	4.6	6.2	5.0	6.0	5.5	6.1
U.S. (Median of the States)	6.1	6.6	5.0	5.1	5.1	5.2	4.9	4.9	4.9	5.8	5.9
** More than 2 drinks per day for males											
** More than 1 drink per day for females											
Tobacco Smoking in Past 30 Days (%)											
Missouri	23.9	25.0	21.1	23.1	24.9	24.5	23.2	23.4	24.1	27.2	26.5
U.S. (Median of the States)	19.6	21.2	17.3	17.9	18.3	19.8	20.1	20.6	20.9	22.0	23.2
Former Smokers (%)											
Missouri	23.9	25.0	26.1	24.7	25.0	23.4	26.8	24.2	24.2	24.3	25.5
U.S. (Median of the States)	25.2	25.1	25.1	25.5	25.2	24.6	24.7	24.8	24.1	25.2	24.7
Limited in Any Activities Due to Physical, Mental, or Emotional Problems											
Missouri	23.7	28.1	24.8	22.1	25.0	21.4		20.9		19.3	
U.S. (Median of the States)	20.1	24.3	21.2	18.9	20.6	18.9		18.6		18.3	
Depressive Disorder Identified by Health or Mental Health Professional in Lifetime (%)											
Missouri	20.1	20.1									
U.S. (Median of the States)	18.0	17.5									

Values shown are weighted estimates based on sampling. Apparent differences in values might not be statistically significant.

Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention: Behavioral Risk Factor Surveillance System.

**PREGNANCY RISK ASSESSMENT MONITORING SYSTEM (PRAMS)
TABLES FOR ALCOHOL, TOBACCO, AND MENTAL HEALTH**

Survey results of samples of Missouri women after having a live birth:

	2010	2009	2007
ALCOHOL			
Binge Alcohol Use (4 or More Drinks on One Occasion) During 3 Months Before Pregnancy (%)	28.0	30.7	
Any Alcohol Use During 3 Months Before Pregnancy (%)	57.7	58.6	57.7
Any Alcohol Use During Last 3 Months of Pregnancy (%)	5.2	4.8	4.5
Change in Drinking from 3 Months Before Pregnancy to Last 3 Months of Pregnancy (%)			
Non-Drinkers	42.1	41.4	42.3
Drinkers	57.9	58.6	57.7
Drinkers Who Quit, Among Total Sample	52.7	53.8	53.2
Drinkers who Reduced Number of Drinks	3.8	2.1	2.7
Drinkers who Increased or Maintained Same Number of Drinks	1.2	2.6	1.8
Non-Drinkers Who Began or Resumed Drinking	0.2	0.0	0.0
Drinkers who Quit During Pregnancy, Among Drinkers Only (%) *	91.0	91.8	92.2
TOBACCO			
Tobacco Smoking During Three Months Before Pregnancy (%)	33.5	35.2	31.7
Tobacco Smoking During Last Three Months of Pregnancy (%)	16.7	18.9	18.4
Change in Smoking from 3 Months Before Pregnancy to Last 3 Months of Pregnancy (%)			
Non-Smokers	66.7	64.6	68.3
Smokers	33.3	35.4	31.7
Smokers Who Quit, Among Total Sample	16.6	16.4	13.4
Smokers Who Reduced Number of Cigarettes Per Day	11.1	11.1	12.2
Smokers Who Increased or Maintained Same Number of Cigarettes Per Day	5.7	7.7	6.1
Non-Smokers Who Began or Resumed Smoking	0.0	0.2	0.0
Tobacco Smokers Who Quit During Pregnancy, Among Smokers Only (%) *	49.7	46.6	42.1
MENTAL HEALTH			
Anxiety During 3 Months Before Pregnancy (%)	13.4	13.1	
Depression During 3 Months Before Pregnancy (%)	13.1	13.6	

Data Source:

U.S. Department of Health and Human Services, Centers for Disease Control and Prevention: Pregnancy Risk Assessment Monitoring System.

Accessed from: <http://apps.nccd.cdc.gov/cPONDER/default.aspx>

* Indicates calculated values

Blank data cells indicate data not collected or available.

MONITORING THE FUTURE SURVEY: TABLES FOR GRADE 8

Estimated Daily Substance Use in Past 30 Days (%): U.S. Students in Grade 8

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	0.3	0.4	0.5	0.5	0.7	0.6	0.5	0.5	0.6	0.8	0.7	0.9
Been Drunk Daily	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Cigarettes	1.9	2.4	2.9	2.7	3.1	3.0	4.0	4.0	4.4	4.5	5.1	5.5
Smokeless Tobacco	0.5	0.8	0.9	0.8	0.8	0.8	0.7	0.7	1.0	0.8	0.8	1.2
Marijuana/Hashish	1.1	1.3	1.2	1.0	0.9	0.8	1.0	1.0	0.8	1.0	1.2	1.3

Estimated Past-30 Day Substance Use (%): U.S. Students in Grade 8

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	11	12.7	13.8	14.9	15.9	15.9	17.2	17.1	18.6	19.7	19.6	21.5
Been Drunk (past 30 days)	3.6	4.4	5.0	5.4	5.4	5.5	6.2	6.0	6.2	6.7	6.7	7.7
Binge Alcohol Use (past 2 weeks)*	5.1	6.4	7.2	7.8	8.1	8.3	8.7	8.4	9.4	9.8	10.3	11.0
Cigarettes	4.9	6.1	7.1	6.5	6.8	7.1	8.7	9.3	9.2	10.2	10.7	12.2
Smokeless Tobacco	2.8	3.5	4.1	3.7	3.5	3.2	3.7	3.3	4.1	4.1	3.3	4.0
Any Illicit Drug	7.7	8.5	9.5	8.1	7.6	7.4	8.1	8.5	8.4	9.7	10.4	11.7
Marijuana/Hashish	6.5	7.2	8.0	6.5	5.8	5.7	6.5	6.6	6.4	7.5	8.3	9.2
Inhalants	2.7	3.2	3.6	3.8	4.1	3.9	4.1	4.2	4.5	4.1	3.8	4.0
Hallucinogens	0.6	1.0	1.0	0.9	0.9	1.0	0.9	1.1	1.0	1.2	1.2	1.6
LSD	0.3	0.5	0.6	0.5	0.5	0.5	0.4	0.5	0.5	0.6	0.7	1.0
MDMA "Ecstasy"	0.5	0.6	1.1	0.6	0.8	0.6	0.7	0.6	0.8	0.7	1.4	1.8
Cocaine (all forms)	0.5	0.8	0.6	0.8	0.8	0.9	1.0	1.0	0.9	0.9	1.1	1.2
Crack Cocaine	0.3	0.5	0.4	0.5	0.5	0.6	0.6	0.6	0.6	0.7	0.8	0.8
Amphetamines	1.3	1.8	1.8	1.9	2.2	2.0	2.1	2.3	2.3	2.7	2.8	3.2
Methamphetamine	0.5	0.4	0.7	0.5	0.7	0.6	0.6	0.7	0.6	1.2	1.1	1.3
Heroin	0.2	0.4	0.4	0.4	0.4	0.4	0.3	0.5	0.5	0.4	0.5	0.6
Tranquilizers	0.8	1.0	1.2	1.2	1.2	1.1	1.3	1.3	1.2	1.4	1.2	1.2
Rohypnol	0.1	0.6	0.2	0.2	0.1	0.3	0.4	0.2	0.2	0.1	0.2	0.4
Steroids	0.3	0.4	0.3	0.4	0.5	0.4	0.5	0.5	0.5	0.7	0.8	0.7
Injection Drug Use	0.2	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.4

Estimated Lifetime Substance Use (%): U.S. Students in Grade 8

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	29.5	33.1	35.8	36.6	38.9	38.9	40.5	41.0	43.9	45.6	47.0	50.5
Ever Been Drunk	12.8	14.8	16.3	17.4	18.0	17.9	19.5	19.5	19.9	20.3	21.3	23.4
Cigarettes	15.5	18.4	20.0	20.1	20.5	22.1	24.6	25.9	27.9	28.4	31.4	36.6
Smokeless Tobacco	8.1	9.7	9.9	9.6	9.8	9.1	10.2	10.1	11.0	11.3	11.2	11.7
Any Illicit Drug	18.5	20.1	21.4	19.9	19.6	19.0	20.9	21.4	21.5	22.8	24.5	26.8
Marijuana/Hashish	15.2	16.4	17.3	15.7	14.6	14.2	15.7	16.5	16.3	17.5	19.2	20.4
Inhalants	11.8	13.1	14.5	14.9	15.7	15.6	16.1	17.1	17.3	15.8	15.2	17.1
Hallucinogens	2.8	3.3	3.4	3.0	3.3	3.1	3.4	3.8	3.5	4.0	4.1	5.2
LSD	1.3	1.7	1.8	1.7	1.9	1.6	1.6	1.9	1.8	2.1	2.5	3.4
MDMA "Ecstasy"	2.0	2.6	3.3	2.2	2.4	2.3	2.5	2.8	2.8	3.2	4.3	5.2
Cocaine (all forms)	1.9	2.2	2.6	2.6	3.0	3.1	3.4	3.7	3.4	3.6	3.6	4.3
Crack Cocaine	1.0	1.5	1.5	1.7	2.0	2.1	2.3	2.4	2.4	2.5	2.5	3.0
Amphetamines	4.5	5.2	5.7	6.0	6.8	6.5	7.3	7.4	7.5	8.4	8.7	10.2
Methamphetamine	1.3	1.3	1.8	1.6	2.3	1.8	2.7	3.1	2.5	3.9	3.5	4.4
Heroin	0.8	1.2	1.3	1.3	1.4	1.3	1.4	1.5	1.6	1.6	1.6	1.7
Tranquilizers	3.0	3.4	4.4	3.9	3.9	3.9	4.3	4.1	4.0	4.4	4.3	5.0
Rohypnol	1.0	2.0	0.9	0.7	0.7	1.0	1.0	1.1	1.0	1.0	0.8	1.1
Steroids	1.2	1.2	1.1	1.3	1.4	1.5	1.6	1.7	1.9	2.5	2.5	2.8
Injection Drug Use	0.6	0.8	0.9	0.9	0.9	0.9	1.0	1.0	1.1	1.0	1.0	1.2

* Binge alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks.

Data Source: Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2013).

Monitoring the Future national survey results on drug use, 1975-2012: Volume 1, Secondary school students.

Ann Arbor: Institute for Social Research, The University of Michigan.

MONITORING THE FUTURE SURVEY: TABLES FOR GRADE 10

Estimated Daily Substance Use in Past 30 Days (%): U.S. Students in Grade 10

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	1.0	0.8	1.1	1.1	1.0	1.4	1.4	1.3	1.3	1.5	1.8	1.9
Been Drunk Daily	0.4	0.2	0.3	0.4	0.3	0.5	0.5	0.4	0.4	0.5	0.5	0.6
Cigarettes	5.0	5.5	6.6	6.3	5.9	7.2	7.6	7.5	8.3	8.9	10.1	12.2
Smokeless Tobacco	2.0	1.7	2.5	1.9	1.4	1.6	1.7	1.9	1.6	1.8	1.7	2.2
Marijuana/Hashish	3.5	3.6	3.3	2.8	2.7	2.8	2.8	3.1	3.2	3.6	3.9	4.5

Estimated Past-30 Day Substance Use (%): U.S. Students in Grade 10

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	27.6	27.2	28.9	30.4	28.8	33.4	33.8	33.2	35.2	35.4	35.4	39.0
Been Drunk (past 30 days)	14.5	13.7	14.7	15.5	14.4	18.1	18.8	17.6	18.5	18.2	18.3	21.9
Binge Alcohol Use (past 2 weeks)*	15.6	14.7	16.3	17.5	16.0	19.6	19.9	19.0	19.9	20.0	20.3	22.8
Cigarettes	10.8	11.8	13.6	13.1	12.3	14.0	14.5	14.9	16.0	16.7	17.7	21.3
Smokeless Tobacco	6.4	6.6	7.5	6.5	5.0	6.1	5.7	5.6	4.9	5.3	6.1	6.9
Any Illicit Drug	18.6	19.2	18.5	17.8	15.8	16.9	16.8	17.3	18.3	19.5	20.8	22.7
Marijuana/Hashish	17.0	17.6	16.7	15.9	13.8	14.2	14.2	15.2	15.9	17.0	17.8	19.8
Inhalants	1.4	1.7	2.0	2.2	2.1	2.5	2.3	2.2	2.4	2.2	2.4	2.4
Hallucinogens	1.2	1.4	1.6	1.4	1.3	1.7	1.5	1.5	1.6	1.5	1.6	2.1
LSD	0.5	0.7	0.7	0.5	0.7	0.7	0.7	0.6	0.6	0.6	0.7	1.5
MDMA "Ecstasy"	1.0	1.6	1.9	1.3	1.1	1.2	1.2	1.0	0.8	1.1	1.8	2.6
Cocaine (all forms)	0.8	0.7	0.9	0.9	1.2	1.3	1.5	1.5	1.7	1.3	1.6	1.3
Crack Cocaine	0.4	0.4	0.5	0.4	0.5	0.5	0.7	0.7	0.8	0.7	1.0	0.7
Heroin	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.5	0.3
Amphetamines	2.8	3.1	3.3	3.3	2.8	4.0	3.5	3.7	4.0	4.3	5.2	5.6
Methamphetamine	0.6	0.5	0.7	0.6	0.7	0.4	0.7	1.1	1.3	1.4	1.8	1.5
Tranquilizers	1.7	1.9	2.2	2.0	1.9	2.6	2.4	2.3	2.3	2.4	2.9	2.9
Rohypnol	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.2	0.4	0.2
Steroids	0.4	0.5	0.5	0.5	0.5	0.5	0.6	0.6	0.8	0.8	1.0	0.9
Injection Drug Use	0.2	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.4

Estimated Lifetime Substance Use (%): U.S. Students in Grade 10

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	54.0	56.0	58.2	59.1	58.3	61.7	61.5	63.2	64.2	66.0	66.9	70.1
Ever Been Drunk	34.6	35.9	36.9	38.6	37.2	41.2	41.4	42.1	42.3	42.4	44.0	48.2
Cigarettes	27.7	30.4	33.0	32.7	31.7	34.6	36.1	38.9	40.7	43.0	47.4	52.8
Smokeless Tobacco	15.4	15.6	16.8	15.2	12.2	15.1	15.0	14.5	13.8	14.6	16.9	19.5
Any Illicit Drug	36.8	37.7	37.0	36.0	34.1	35.6	36.1	38.2	39.8	41.4	44.6	45.6
Marijuana/Hashish	33.8	34.5	33.4	32.3	29.9	31.0	31.8	34.1	35.1	36.4	38.7	40.1
Inhalants	9.9	10.1	12.0	12.3	12.8	13.6	13.3	13.1	12.4	12.7	13.5	15.2
Hallucinogens	5.2	6.0	6.1	6.1	5.5	6.4	6.1	5.8	6.4	6.9	7.8	8.9
LSD	2.6	2.8	3.0	3.0	2.6	3.0	2.7	2.5	2.8	3.5	5.0	6.3
MDMA "Ecstasy"	5.0	6.6	6.4	5.5	4.3	5.2	4.5	4.0	4.3	5.4	6.6	8.0
Cocaine (all forms)	3.3	3.3	3.7	4.6	4.5	5.3	4.8	5.2	5.4	5.1	6.1	5.7
Crack Cocaine	1.4	1.6	1.8	2.1	2.0	2.3	2.2	2.5	2.6	2.7	3.6	3.1
Amphetamines	8.9	9.0	10.6	10.3	9.0	11.1	11.2	11.1	11.9	13.1	14.9	16.0
Methamphetamine	1.8	2.1	2.5	2.8	2.4	2.8	3.2	4.1	5.3	5.2	6.1	6.4
Heroin	1.1	1.2	1.3	1.5	1.2	1.5	1.4	1.5	1.5	1.5	1.8	1.7
Tranquilizers	6.3	6.8	7.3	7.0	6.8	7.4	7.2	7.1	7.3	7.8	8.8	9.2
Rohypnol	0.8	1.2	1.4	0.7	0.9	1.3	0.8	1.0	1.2	1.0	1.3	1.5
Steroids	1.3	1.4	1.6	1.3	1.4	1.8	1.8	2.0	2.4	3.0	3.5	3.5
Injection Drug Use	0.7	0.8	0.8	0.9	0.7	0.9	0.9	0.8	0.8	0.9	1.0	0.8

* Binge alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks.

Data Source: Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2013). *Monitoring the Future national survey results on drug use, 1975-2012: Volume 1, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan.

MONITORING THE FUTURE SURVEY: TABLES FOR GRADE 12

Estimated Daily Substance Use in Past 30 Days (%): U.S. Students in Grade 12

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	2.5	2.1	2.7	2.5	2.8	3.1	3.0	3.1	2.8	3.2	3.5	3.6
Been Drunk Daily	1.5	1.3	1.6	1.1	1.4	1.3	1.6	1.5	1.8	1.6	1.2	1.4
Cigarettes	9.3	10.3	10.7	11.2	11.4	12.3	12.2	13.6	15.6	15.8	16.9	19.0
Smokeless Tobacco	3.2	3.1	3.1	2.9	2.7	2.8	2.2	2.5	2.8	2.2	2.0	2.8
Marijuana/Hashish	6.5	6.6	6.1	5.2	5.4	5.1	5.0	5.0	5.6	6.0	6.0	5.8

Estimated Past-30 Day Substance Use (%): U.S. Students in Grade 12

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	41.5	40.0	41.2	43.5	43.1	44.4	45.3	47.0	48.0	47.5	48.6	49.8
Been Drunk (past 30 days)	28.1	25.0	26.8	27.4	27.6	28.7	30.0	30.2	32.5	30.9	30.3	32.7
Binge Alcohol Use (past 2 weeks)	23.7	21.6	23.2	25.2	24.6	25.9	25.4	27.1	29.2	27.9	28.6	29.7
Cigarettes	17.1	18.7	19.2	20.1	20.4	21.6	21.6	23.2	25.0	24.4	26.7	29.5
Smokeless Tobacco	7.9	8.3	8.5	8.4	6.5	6.6	6.1	7.6	6.7	6.7	6.5	7.8
Any Illicit Drug	25.2	25.2	23.8	23.3	22.3	21.9	21.5	23.1	23.4	24.1	25.4	25.7
Marijuana/Hashish	22.9	22.6	21.4	20.6	19.4	18.8	18.3	19.8	19.9	21.2	21.5	22.4
Inhalants	0.9	1.0	1.4	1.2	1.4	1.2	1.5	2.0	1.5	1.5	1.5	1.7
Hallucinogens	1.6	1.6	1.9	1.6	2.2	1.7	1.5	1.9	1.9	1.8	2.3	3.3
LSD	0.8	0.8	0.8	0.5	1.1	0.6	0.6	0.7	0.7	0.6	0.7	2.3
PCP	0.5	0.8	0.8	0.5	0.6	0.5	0.4	0.7	0.4	0.6	0.4	0.5
MDMA "Ecstasy"	0.9	2.3	1.4	1.8	1.8	1.6	1.3	1.0	1.2	1.3	2.4	2.8
Cocaine (all forms)	1.1	1.1	1.3	1.3	1.9	2.0	2.5	2.3	2.3	2.1	2.3	2.1
Crack Cocaine	0.6	0.5	0.7	0.6	0.8	0.9	0.9	1.0	1.0	0.9	1.2	1.1
Amphetamines	3.3	3.7	3.3	3.0	2.9	3.7	3.7	3.9	4.6	5.0	5.5	5.6
Methamphetamine	0.5	0.6	0.5	0.5	0.6	0.6	0.9	0.9	1.4	1.7	1.7	1.5
Crystal Methamphetamine "Ice"	0.4	0.6	0.6	0.5	0.6	0.6	0.7	0.9	0.8	0.8	1.2	1.1
Heroin	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.4	0.5	0.4
Other Narcotics	3.0	3.6	3.6	4.1	3.8	3.8	3.8	3.9	4.3	4.1	4.0	3.0
Barbiturates	2.0	1.8	2.2	2.5	2.8	2.7	3.0	3.3	2.9	2.9	3.2	2.8
Methaqualone	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.5	0.5	0.4	0.3	0.5
Tranquilizers	2.1	2.3	2.5	2.7	2.6	2.6	2.7	2.9	3.1	2.8	3.3	2.9
Steroids	0.9	0.7	1.1	1.0	1.0	1.0	1.1	0.9	1.6	1.3	1.4	1.3
Injection Drug Use	0.3	0.4	0.4	0.1	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2

Estimated Lifetime Substance Use (%): U.S. Students in Grade 12

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	69.4	70.0	71.0	72.3	71.9	72.2	72.7	75.1	76.8	76.6	78.4	79.7
Ever Been Drunk	54.2	51.0	54.1	56.5	54.7	55.1	56.4	57.5	60.3	58.1	61.6	63.9
Cigarettes	39.5	40.0	42.2	43.6	44.7	46.2	47.1	50.0	52.8	53.7	57.2	61.0
Smokeless Tobacco	17.4	16.9	17.6	16.3	15.6	15.1	15.2	17.5	16.7	17.0	18.3	19.7
Any Illicit Drug	49.1	49.9	48.2	46.7	47.4	46.8	48.2	50.4	51.1	51.1	53.0	53.9
Marijuana/Hashish	45.2	45.5	43.8	42.0	42.6	41.8	42.3	44.8	45.7	46.1	47.8	49.0
Inhalants	7.9	8.1	9.0	9.5	9.9	10.5	11.1	11.4	10.9	11.2	11.7	13.0
Hallucinogens	7.5	8.3	8.6	7.4	8.7	8.4	8.3	8.8	9.7	10.6	12.0	14.7
LSD	3.8	4.0	4.0	3.1	4.0	3.4	3.3	3.5	4.6	5.9	8.4	10.9
PCP	1.6	2.3	1.8	1.7	1.8	2.1	2.2	2.4	1.6	2.5	3.1	3.5
MDMA "Ecstasy"	7.2	8.0	7.3	6.5	6.2	6.5	6.5	5.4	7.5	8.3	10.5	11.7
Cocaine (all forms)	4.9	5.2	5.5	6.0	7.2	7.8	8.5	8.0	8.1	7.7	7.8	8.2
Crack Cocaine	2.1	1.9	2.4	2.4	2.8	3.2	3.5	3.5	3.9	3.6	3.8	3.7
Amphetamines	12.0	12.2	11.1	9.9	10.5	11.4	12.4	13.1	15.0	14.4	16.8	16.2
Methamphetamine	1.7	2.1	2.3	2.4	2.8	3.0	4.4	4.5	6.2	6.2	6.7	6.9
Crystal Methamphetamine "Ice"	1.7	2.1	1.8	2.1	2.8	3.4	3.4	4.0	4.0	3.9	4.7	4.1
Heroin	1.1	1.4	1.6	1.2	1.3	1.5	1.4	1.5	1.5	1.5	1.7	1.8
Other Narcotics	12.2	13.0	13.0	13.2	13.2	13.1	13.4	12.8	13.5	13.2	13.5	9.9
Barbiturates	6.9	7.0	7.5	8.2	8.5	9.3	10.2	10.5	9.9	8.8	9.5	8.7
Methaqualone	0.8	0.6	0.4	0.7	0.8	1.0	1.2	1.3	1.3	1.0	1.5	1.1
Tranquilizers	8.5	8.7	8.5	9.3	8.9	9.5	10.3	9.9	10.6	10.2	11.4	10.3
Steroids	1.8	1.8	2.0	2.2	2.2	2.2	2.7	2.6	3.4	3.5	4.0	3.7
Injection Drug Use	0.7	0.9	1.1	0.6	0.7	0.7	0.8	0.9	0.7	0.7	0.8	0.7

* Binge alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks.

Data Source: Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2013).
Monitoring the Future national survey results on drug use, 1975-2012: Volume 1, Secondary school students.
 Ann Arbor: Institute for Social Research, The University of Michigan.

MONITORING THE FUTURE SURVEY: TABLES FOR COLLEGE STUDENTS

Estimated Daily Substance Use in Past 30 Days (%): U.S. College Students

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	3.9	3.8	3.6	4.3	4.0	4.3	4.8	4.6	3.7	4.3	5.0	4.7
Been Drunk Daily	0.4	1.3	0.3	0.7	0.5	0.7	0.6	0.5	0.8	1.1	0.8	0.5
Cigarettes	5.2	7.3	7.6	8.0	9.2	9.3	9.2	12.4	13.8	13.8	15.9	15.0
Marijuana/Hashish	4.8	4.7	4.4	4.9	3.9	3.5	4.3	4.0	4.5	4.7	4.1	4.5

Estimated Past-30 Day Substance Use (%): U.S. College Students

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	67.7	63.5	65.0	65.8	69.0	66.6	65.4	67.9	67.7	66.2	68.9	67.0
Been Drunk (past 30 days)	40.1	39.9	43.6	42.4	45.3	46.8	47.6	43.1	47.4	40.4	44.4	44.7
Binge Alcohol Use (past 2 weeks)*	37.4	36.1	37.0	36.9	40.0	41.1	40.2	40.1	41.7	38.5	40.1	40.9
Cigarettes	12.5	15.2	16.4	17.9	17.9	19.9	19.2	23.8	24.3	22.5	26.7	25.7
Any Illicit Drug	22.3	21.4	19.2	20.7	18.9	19.3	19.2	19.5	21.2	21.4	21.5	21.9
Marijuana/Hashish	20.5	19.4	17.5	18.5	17.0	16.8	16.7	17.1	18.9	19.3	19.7	20.2
Inhalants	0.2	0.3	0.5	0.1	0.4	0.1	0.4	0.3	0.4	0.4	0.7	0.4
Hallucinogens	1.1	1.2	1.4	1.0	1.7	1.3	0.9	1.2	1.3	1.8	1.2	1.8
LSD	0.4	0.5	0.7	0.3	0.8	0.3	0.3	0.1	0.2	0.2	0.2	1.0
MDMA "Ecstasy"	1.4	0.7	1.0	0.5	0.6	0.4	0.6	0.8	0.7	1.0	0.7	1.5
Cocaine (all forms)	1.1	1.2	1.0	1.3	1.2	1.7	1.8	1.8	2.4	1.9	1.6	1.9
Crack Cocaine	0.0	0.1	0.1	0.1	0.1	0.1	**	0.1	0.4	0.4	0.3	0.1
Amphetamines	4.6	4.5	4.1	3.4	2.8	3.1	2.5	2.9	3.2	3.1	3.0	3.3
Methamphetamine	0.0	0.0	0.0	0.1	0.0	0.1	0.2	0.1	0.2	0.6	0.2	0.5
Crystal Methamphetamine "Ice"	0.3	0.0	0.2	0.0	0.0	0.1	**	0.2	0.1	0.3	0.0	0.1
Heroin	0.1	0.0	0.0	0.1	**	0.1	0.2	0.1	0.1	0.0	0.0	0.1
Other Narcotics	2.2	2.1	2.3	2.7	2.3	2.2	3.1	3.1	3.0	2.3	1.6	1.7
Barbiturates	0.8	0.8	0.6	1.2	1.4	1.4	1.3	1.3	1.5	1.7	1.7	1.5
Tranquilizers	1.1	1.6	1.3	2.2	1.6	1.8	2.1	2.2	2.7	2.8	3.0	1.5
Steroids	0.0	0.2	0.0	0.2	**	0.1	**	**	**	0.1	**	0.3
Injection Drug Use	0.2	0.0	0.0	0.1	0.0	**	0.1	0.1	0.1	0.1	**	**

Estimated Lifetime Substance Use (%): U.S. College Students

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	81.0	80.5	82.3	82.6	85.3	83.1	84.7	86.6	84.6	86.2	86.0	86.1
Ever Been Drunk	70.0	67.9	70.5	69.1	72.5	71.6	73.1	72.9	73.4	74.9	75.1	76.1
Any Illicit Drug	50.5	49.2	49.1	51.4	49.5	50.5	50.6	52.3	52.2	53.9	51.8	53.6
Marijuana/Hashish	49.1	46.6	46.8	47.5	46.8	47.5	46.9	49.1	49.1	50.7	49.5	51.0
Inhalants	5.7	3.7	5.5	6.9	4.9	6.3	7.4	7.1	8.5	9.7	7.7	9.6
Hallucinogens	7.6	7.4	7.8	8.0	8.5	9.1	10.6	11.0	12.0	14.5	13.6	14.8
LSD	3.1	3.7	4.0	3.3	4.3	3.3	3.5	3.7	5.6	8.7	8.6	12.2
MDMA "Ecstasy"	8.7	6.8	6.2	6.5	6.2	5.4	6.9	8.3	10.2	12.9	12.7	14.7
Cocaine (all forms)	5.2	5.5	6.6	8.1	7.2	8.5	7.7	8.8	9.5	9.2	8.2	8.6
Crack Cocaine	0.7	0.8	1.2	1.0	1.4	1.3	2.3	1.7	2.0	3.1	1.9	2.0
Amphetamines	14.4	13.4	12.1	11.8	9.1	11.2	10.7	12.3	12.7	12.3	11.9	12.4
Methamphetamine	0.3	0.6	1.1	1.0	1.9	1.9	2.9	4.1	5.2	5.8	5.0	5.3
Crystal Methamphetamine "Ice"	0.6	0.2	0.8	0.7	1.1	1.3	1.7	2.4	2.2	2.9	2.0	2.3
Heroin	0.5	0.6	0.7	0.8	0.7	0.5	0.7	0.5	0.9	1.0	1.0	1.2
Other Narcotics	10.3	12.4	12.2	14.0	12.4	14.1	14.6	14.4	13.8	14.2	12.2	11.0
Barbiturates	3.5	3.6	5.3	6.0	6.4	5.9	6.3	8.5	7.2	5.7	5.9	6.0
Tranquilizers	6.4	7.1	8.1	9.2	8.6	9.1	10.0	11.9	10.6	11.0	10.7	9.7
Steroids	0.4	1.1	0.7	1.3	1.6	0.6	1.9	1.0	1.6	1.2	1.2	1.5
Injection Drug Use	0.2	0.3	0.1	0.1	0.0	0.1	0.3	0.3	0.1	0.1	0.3	0.2

* Binge alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks.

** Indicates value is less than 0.05 percent but greater than 0.0 percent.

Data Source: Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2013).

Monitoring the Future national survey results on drug use, 1975-2012: Volume 2, College students and adults ages 19-50.

Ann Arbor: Institute for Social Research, The University of Michigan.

MONITORING THE FUTURE SURVEY: TABLES FOR YOUNG ADULTS AGES 19-28

Estimated Daily Substance Use in Past 30 Days (%): U.S. Young Adults Ages 19-28

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	5.5	5.2	4.6	5.3	5.3	5.6	5.4	5.2	4.5	5.1	4.7	4.4
Been Drunk Daily	0.4	0.7	0.7	1.0	0.5	0.6	0.6	0.5	0.7	0.8	0.6	0.4
Cigarettes	12.8	13.8	14.8	15.0	16.7	17.3	18.6	19.6	20.8	20.3	21.2	21.2
Marijuana/Hashish	5.6	6.1	5.3	5.4	5.1	5.0	5.0	4.9	5.0	5.3	4.5	5.0

Estimated Past-30 Day Substance Use (%): U.S. Young Adults Ages 19-28

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	69.5	68.8	68.4	69.4	68.9	69.5	68.7	68.6	68.4	67.0	68.3	67.0
Been Drunk (past 30 days)	39.1	39.5	39.4	40.5	40.7	41.4	42.1	39.0	39.0	37.8	37.1	36.8
Binge Alcohol Use (past 2 weeks)*	35.5	36.5	35.9	36.7	37.9	37.8	37.6	37.0	37.1	35.8	35.9	35.9
Cigarettes	19.7	21.3	22.4	23.3	24.6	26.2	27.0	28.6	29.2	28.4	29.2	30.2
Any Illicit Drug	19.9	20.6	18.9	19.8	19.3	18.9	18.5	18.6	19.1	19.9	18.9	18.8
Marijuana/Hashish	17.7	18.3	16.1	17.0	16.0	16.0	15.7	15.8	16.5	17.3	16.9	16.7
Inhalants	0.3	0.1	0.1	0.2	0.4	0.2	0.3	0.2	0.3	0.3	0.5	0.4
Hallucinogens	0.6	0.9	1.0	0.8	0.9	0.9	0.7	0.8	0.9	1.2	0.9	1.2
LSD	0.3	0.3	0.4	0.2	0.4	0.2	0.2	0.1	0.1	0.2	0.3	0.7
PCP	0.0	0.1	0.0	**	0.1	**	**	0.0	0.1	0.1	0.1	0.0
MDMA "Ecstasy"	1.0	0.7	0.8	0.6	0.6	0.5	0.7	0.6	0.6	0.8	1.3	1.8
Cocaine (all forms)	1.3	1.5	1.4	1.8	1.9	2.1	2.3	2.2	2.2	2.4	2.2	2.2
Crack Cocaine	0.1	0.2	0.1	0.2	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.4
Amphetamines	3.2	3.0	2.9	2.5	2.2	2.3	2.2	2.1	2.4	2.5	2.5	2.4
Methamphetamine	0.4	0.3	0.2	0.3	0.3	0.6	0.5	0.7	0.6	0.7	1.0	1.0
Crystal Methamphetamine "Ice"	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.6	0.4	0.4	0.5	0.4
Heroin	0.2	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	**	0.3
Other Narcotics	2.7	2.9	3.4	3.2	3.6	3.4	3.2	3.5	3.0	2.9	2.9	1.7
Barbiturates	1.1	1.1	1.1	1.2	1.9	1.6	1.5	1.7	1.8	1.5	1.5	1.7
Tranquilizers	1.9	2.3	2.2	2.8	2.7	2.8	2.3	2.6	2.7	2.4	2.8	2.1
Steroids	0.1	0.2	0.5	0.3	0.2	0.4	0.1	0.1	0.1	0.1	0.1	0.1
Injection Drug Use	0.2	0.2	0.1	0.1	**	**	0.1	0.1	0.1	**	**	0.2

Estimated Lifetime Substance Use (%): U.S. Young Adults Ages 19-28

	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Alcohol	86.5	87.4	87.5	87.9	88.4	87.9	88.9	89.1	89.4	89.3	90.2	89.9
Ever Been Drunk	78.9	78.9	79.0	78.2	80.1	80.1	80.9	79.9	80.1	80.9	81.2	81.1
Any Illicit Drug	58.9	59.1	58.4	59.3	59.3	59.8	59.7	60.4	60.5	60.2	59.0	58.1
Marijuana/Hashish	56.5	56.3	55.9	56.0	55.9	56.7	56.7	57.0	57.4	57.2	56.8	55.7
Inhalants	7.2	7.2	7.9	8.9	9.5	9.1	10.9	10.3	11.6	12.2	12.4	12.8
Hallucinogens	12.2	13.0	13.9	14.2	14.8	16.0	17.2	17.6	19.3	19.7	19.6	18.3
LSD	6.2	6.1	7.2	7.3	8.1	9.6	10.1	11.2	13.4	14.6	15.1	16.0
PCP	1.1	1.7	1.6	1.6	2.2	2.1	2.4	2.0	2.7	3.0	2.5	3.1
MDMA "Ecstasy"	11.4	11.3	12.3	11.5	13.1	13.1	14.4	14.9	16.0	15.3	14.6	13.0
Cocaine (all forms)	11.9	12.5	13.6	13.9	14.8	14.7	15.2	14.3	15.2	14.7	13.5	13.1
Crack Cocaine	2.7	2.9	3.6	3.3	4.3	3.9	4.4	4.1	4.2	4.7	4.3	4.7
Amphetamines	17.4	16.5	16.1	14.9	14.6	15.3	15.6	14.6	15.9	15.2	14.8	15.0
Methamphetamine	3.5	3.2	4.3	4.7	6.3	6.7	7.3	8.3	9.0	8.9	9.1	9.0
Crystal Methamphetamine "Ice"	2.6	3.1	2.8	3.4	3.6	3.7	4.7	4.4	4.7	4.7	4.1	4.0
Heroin	1.6	1.7	1.8	1.6	1.9	1.6	1.9	1.7	1.9	1.9	1.8	2.0
Other Narcotics	17.6	18.2	19.0	18.5	19.5	18.8	18.7	17.8	17.6	16.8	13.9	11.5
Barbiturates	7.2	7.9	8.6	9.5	10.6	9.8	9.5	10.0	9.7	8.7	8.0	7.8
Tranquilizers	13.3	13.8	14.3	13.8	15.8	14.5	15.0	14.5	14.9	13.8	13.4	11.9
Steroids	1.7	1.3	1.7	1.8	1.8	1.7	1.8	1.8	1.9	1.8	1.6	1.4
Injection Drug Use	0.5	0.7	0.8	0.5	0.5	0.5	0.6	0.6	0.4	0.5	0.4	0.6

* Binge alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks.

** Indicates value is less than 0.05 percent but greater than 0.0 percent.


Data Source: Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2013).

Monitoring the Future national survey results on drug use, 1975-2012: Volume 2, College students and adults ages 19-50.

Ann Arbor: Institute for Social Research, The University of Michigan.


ALCOHOL — PAST-MONTH USE: MISSOURI AGE GROUP & GRADE LEVEL 3-SURVEY COMPARISON

Estimated Past-Month Alcohol Use (%): Missouri Age Group Comparisons, 2-Year Averages, 2002-2011




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Alcohol Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey


Reported Past 30-Day Alcohol Use (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


CIGARETTE — PAST-MONTH USE: MISSOURI AGE GROUP & GRADE LEVEL 3-SURVEY COMPARISON

Estimated Past-Month Cigarette Use (%): Missouri Age Group Comparisons, 2-Year Averages, 2002-2011




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Cigarette Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey


Reported Past 30-Day Cigarette Use (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.


MARIJUANA — PAST-MONTH USE: MISSOURI AGE GROUP & GRADE LEVEL 3-SURVEY COMPARISON

Estimated Past-Month Marijuana Use (%): Missouri Age Group Comparisons, 2-Year Averages, 2002-2011




Data Source: U.S. Department of Health and Human Services, SAMHSA. National Survey on Drug Use and Health.

Estimated Past 30-Day Marijuana Use (%): Missouri Students in Grades 9-12, 1995-2009



Data Source: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Youth Risk Behavior Survey

Reported Past 30-Day Marijuana Use (%): Missouri Students in Grades 6-12, 2004-2012



Data Source: Missouri Institute of Mental Health. Missouri Student Survey. 2004-2008 data are unweighted; 2010 and 2012 data are weighted.